

An agronomic assessment of greenhouse gas emissions from major cereal crops

BRUCE LINQUIST*, KEES JAN VAN GROENIGEN†‡, MARIA ARLENE ADVIENTO-BORBE*, CAMERON PITTELKOW* and CHRIS VAN KESSEL*

*Department of Plant Sciences, University of California, Davis, CA 95616, USA, †Department of Biological Sciences and Merriam-Powell Center for Environmental Research, Northern Arizona University, Flagstaff, AZ 86011, USA, ‡Department of Botany, School of Natural Sciences, Trinity College Dublin, Dublin 2, Ireland

Abstract

Agricultural greenhouse gas (GHG) emissions contribute approximately 12% to total global anthropogenic GHG emissions. Cereals (rice, wheat, and maize) are the largest source of human calories, and it is estimated that world cereal production must increase by 1.3% annually to 2025 to meet growing demand. Sustainable intensification of cereal production systems will require maintaining high yields while reducing environmental costs. We conducted a meta-analysis (57 published studies consisting of 62 study sites and 328 observations) to test the hypothesis that the global warming potential (GWP) of CH₄ and N₂O emissions from rice, wheat, and maize, when expressed per ton of grain (yield-scaled GWP), is similar, and that the lowest value for each cereal is achieved at near optimal yields. Results show that the GWP of CH₄ and N₂O emissions from rice (3757 kg CO₂ eq ha⁻¹ season⁻¹) was higher than wheat (662 kg CO₂ eq ha⁻¹ season⁻¹) and maize (1399 kg CO₂ eq ha⁻¹ season⁻¹). The yield-scaled GWP of rice was about four times higher (657 kg CO₂ eq Mg⁻¹) than wheat (166 kg CO₂ eq Mg⁻¹) and maize (185 kg CO₂ eq Mg⁻¹). Across cereals, the lowest yield-scaled GWP values were achieved at 92% of maximal yield and were about twice as high for rice (279 kg CO₂ eq Mg⁻¹) than wheat (102 kg CO₂ eq Mg⁻¹) or maize (140 kg CO₂ eq Mg⁻¹), suggesting greater mitigation opportunities for rice systems. In rice, wheat and maize, 0.68%, 1.21%, and 1.06% of N applied was emitted as N₂O, respectively. In rice systems, there was no correlation between CH₄ emissions and N rate. In addition, when evaluating issues related to food security and environmental sustainability, other factors including cultural significance, the provisioning of ecosystem services, and human health and well-being must also be considered.

Keywords: CH₄, corn, global warming potential, grain yield, maize, meta-analysis, methane, N₂O, nitrogen rate, nitrous oxide, rice, wheat, yield-scaled

Received 14 April 2011 and accepted 29 May 2011

Introduction

Global cropland area is roughly 1.5 billion ha (Thenkabail *et al.*, 2010) and the major cereals (rice, wheat, and maize) are produced on approximately 546 million ha (Table 1), representing 36% of this area. These crops provide close to 60% of all human calories, either directly as human food or indirectly as livestock feed (Cassman *et al.*, 2003). The Green Revolution and the corresponding intensification of rice (*Oryza sativa*), wheat (*Triticum aestivum*), and maize (*Zea mays*) systems has largely been responsible for averting a short fall in food supply during previous decades (Cassman, 1999; Tilman, 1999; Burney *et al.*, 2010). With an expanding world population, the demand for these crops will continue to increase at about 1.3% annually to 2025 (Cassman *et al.*, 2003). Broadly, two options exist for increasing cereal production. First, agriculture

can be expanded to new areas that are currently not used for food production. Land for such expansion is available especially in Africa and South America (Deininger & Byerlee, 2011). Second, intensification of existing agricultural land can occur by achieving higher yield per unit of land area (Burney *et al.*, 2010; Godfray *et al.*, 2011), as was achieved in many countries during the Green Revolution. However, because both these options can have negative environmental outcomes, the potential remains for agriculture to further degrade ecosystems in the future. Expansion into new areas can particularly have adverse effects on habitat and biodiversity, whereas agricultural intensification can lead to non-point source pollution and increased greenhouse gas (GHG) emissions (Matson *et al.*, 1997; Vitousek *et al.*, 1997; Tilman, 1999). Proponents of the latter approach (e.g. Godfray *et al.*, 2011) suggest that sustainable intensification, by which higher yields are achieved with no or reduced damage to the environment, will be necessary to meet dual goals of protecting natural resources while ensuring global food security.

Correspondence: Bruce Linquist, tel. (530) 752-3125, fax (530) 752-4361, e-mail: balinquist@ucdavis.edu

Table 1 Area in production and average yields of the world's major cereal crops for 2009 (<http://www.faostat.fao.org>). The "study average yields" are the average yields for each crop in our meta-analysis

Crop	Total production (Mg)	Harvested area (ha)	Average yields (Mg ha ⁻¹)	Study average yields (Mg ha ⁻¹)
Rice (paddy)	678 688 289	161 420 743	4.2	6.05
Wheat	681 915 838	225 437 694	3.0	4.78
Maize	817 110 509	159 531 007	5.1	8.01

It is estimated that agriculture accounts for 10–12% of total global anthropogenic emissions of GHG, which amounts to 60% and 50% of global N₂O and CH₄ emissions, respectively (Smith *et al.*, 2007). N₂O is a more potent GHG than CH₄ with a radiative forcing potential that is approximately 12 times larger (IPCC, 2001). Wheat, maize, and other upland crops are primarily a source of N₂O emissions, with these emissions largely driven by the amount of fertilizer N applied (Cole *et al.*, 1997; van Groenigen *et al.*, 2010). Aerobic upland soils contribute little to CH₄ emissions and may even be a sink for CH₄ in some cases (e.g. Adviento-Borbe *et al.*, 2007). Rice systems are fundamentally different, as rice is typically grown in flooded soils. CH₄ is the dominant GHG produced and emitted in these systems, with emissions being largely controlled by water and residue management practices (Yagi *et al.*, 1997; Wassmann *et al.*, 2000). However, rice systems also emit N₂O, and it has been shown that the intensity of emissions is related to N fertilizer rate (Zou *et al.*, 2007). In rice systems, there is often an inverse relationship between CH₄ and N₂O emissions (Hou *et al.*, 2000). For example, N₂O emissions tend to increase when management practices are implemented to reduce CH₄ emissions, through the use of mid-season drains (Cai *et al.*, 1997; Zou *et al.*, 2007). Rice systems are also unique from other systems in that the majority of CH₄, as well as some N₂O, are emitted through the plant rather than the soil (Yu *et al.*, 1997).

van Groenigen *et al.* (2010) postulated that in a world with increasing food demand and limited land area for expansion of agriculture, N₂O emissions [or global warming potential (GWP)] should be assessed as a function of crop yield (i.e. N₂O produced per unit of grain yield – termed yield-scaled GWP), rather than as a function of area, as is often reported. Ideally, strategies should be identified that allow for the lowest yield-scaled GWP. As GHG emissions are largely driven by fertilizer additions (which tend to increase yield), promoting management practices with low GWP per unit of land area can lead to lower yields. Recently, field study (Ma *et al.*, 2010a; Hoben *et al.*, 2011) and meta-analysis results (van Groenigen *et al.*, 2010) have shown that yield-scaled N₂O emissions were

lowest in intensive crop production systems, where crops were grown close to their yield potential with high N use efficiency. These studies reported that a significant increase in yield-scaled emissions only occurred at high or excessive N rates.

Not only is fertilizer N a major driver of N₂O emissions, it is often the limiting nutrient for crop production and therefore is a major driver of crop yields. Although higher yields can often be obtained with greater fertilizer N inputs, the question is whether the yield increase is large enough to offset the corresponding increase in N₂O emissions and result in an overall lower yield-scaled GWP. In rice systems, the relationship between fertilizer rate and GWP is potentially more complex, as CH₄ emissions are not as closely linked to N fertilizer inputs as N₂O emissions. Although rice systems have been identified as a substantial source of CH₄ emissions, the radiative forcing potential of CH₄ is only 8% of N₂O. Therefore, we tested the hypotheses that (i) yield-scaled GWP estimates are similar for rice, wheat, and maize and (ii) for each cereal the lowest yield-scaled GWP value is achieved at near optimal yield conditions.

Materials and methods

Data

An exhaustive literature survey of peer-reviewed publications was carried out using ISI-Web of Science and Google Scholar (Google Inc., Mountain View, CA, USA) for articles published before January 2011. The literature survey focused on GHG emissions from rice, wheat, and maize systems. Studies had to meet specific criteria to be included in the dataset. First, GHG fluxes must have been measured under field conditions for an entire season. A season included the period from planting to harvest. Several exceptions were made when the portion of the season not included in the study was assumed to contribute only a small percentage to overall seasonal emissions (Majumdar *et al.*, 2002; Parkin & Kaspar, 2006; Datta *et al.*, 2009). All wheat studies were conducted on winter wheat except one (Malhi & Lemke, 2007). In temperate climates, winter wheat is planted in the fall and harvested in the summer, covering almost a full year of GHG emissions.

Second, studies on wheat and maize had to report N₂O emissions and studies on rice N₂O and CH₄ emissions. Although wheat and maize are potential sources or sinks of CH₄, only a few studies (three for wheat and five for maize) measured CH₄ emissions (Table 2), all of them determining that the contribution of CH₄ to the GWP was minor in these systems (e.g. Halvorson *et al.*, 2010). Although soil CO₂ fluxes also represent a source of GHG emissions, on a global scale, they are largely offset by high rates of net primary productivity and atmospheric CO₂ fixation by crop plants, and are therefore estimated to contribute <1% to the GWP of agriculture (Smith *et al.*, 2007). Therefore, CO₂ as a contributor to GWP was not included in our analysis. Third, only studies that reported crop yields were included. In some cases, grain yield data were obtained from other publications or via personal communication (Table 2).

The GWP of N₂O and CH₄ emissions was calculated in units of CO₂ equivalents (CO₂ eq) over a 100-year time horizon. A radiative forcing potential relative to CO₂ of 298 was used for N₂O and 25 for CH₄ (IPCC, 2001). We calculated the combined GWP for N₂O and CH₄ emissions for each individual rice study prior to meta-analysis. To calculate this value for wheat and maize, we added the average GWP for N₂O emissions across studies to the average GWP of CH₄ fluxes.

For each study treatment, the amount of N added as inorganic fertilizer, manure, or green manure was determined. We did not include the amount of N in the previous crop residues as the majority of studies did not report on how it was managed. Also, we did not include N in crop residues applied during the study, as it was not an external input. We divided the studies into four categories based on N fertilization rates (0–<50, 50–<125, 125–200, and >200 kg N ha⁻¹ yr⁻¹) to examine the relationship between N addition and GWP.

The final data set consisted of 16 (17 sites, 116 observations), 19 (20 sites, 122 observations), and 22 (25 sites, 88 observations) studies for rice, maize, and wheat, respectively (Table 2). All of the rice studies were from Asia. For maize, the majority of studies were from the North America (12), four were from Asia, and three from Europe. For wheat, 12 studies were from Asia, six from Europe, and four from North America. There were no studies from Africa, South and Central America, the Middle East, or Australia.

Data analysis

For every study, the net seasonal GHG flux for each individual treatment combination was included as a separate data point (observation) in our meta-analysis. To avoid bias toward multi-year studies, observations were averaged over years when experiments were repeated over time.

We performed meta-analyses using a non-parametric weighting function and generated confidence intervals (CIs) on flux measurements using bootstrapping. Studies were weighted by replication and sampling frequency. When multiple observations were extracted from the same experimental site within the same study (i.e. when GHG fluxes were measured for multiple treatment combinations), we adjusted the weights by the total number of observations from that site:

$$w_i = n \times f / o, \quad (1)$$

where w_i is the weight for observations from the i^{th} site and n is the number of field replicates (i.e., plots per treatment combination). The variable ' f ' is a measure of sampling frequency, and is equal to the number of flux measurements per month. To prevent studies with high temporal sampling frequency from being assigned extreme weights, all studies that measured GHG fluxes more than once a week were assigned the maximum value of $f = 5$. Finally, ' o ' is the total number of flux observations from the i^{th} site. By favoring field experiments that were well replicated and frequently sampled, our weighting approach assigned more weight to more precise flux estimates. Furthermore, our approach ensured that all flux measurements could be included in our analyses without any study dominating the data set. Mean GHG fluxes were estimated as:

$$\bar{U} = \Sigma(U_i \times w_i) / \Sigma(w_i) \quad (2)$$

with U_i as the observation of N₂O or CH₄ flux from the i^{th} site, and w_i as before. Mean yields and yield-scaled GWP were calculated using the same approach. We used METAWIN 2.1 to generate these mean flux sizes and 95% bootstrapped CIs (4999 iterations) (Rosenberg *et al.*, 2000). Mean fluxes for categories of studies (i.e. the three types of cereal crop, and the categories based on N fertilization rate within each crop type) were considered significantly different if their 95% CIs did not overlap.

To assess the potential for reducing yield-scaled GWP, we identified the treatment with the lowest yield-scaled GWP for each site, and repeated all meta-analyses on the reduced data sets.

We tested whether net seasonal GHG fluxes were correlated with N rate using a simple unweighted regression analysis in SPSS v. 19 (SPSS Inc., Chicago, IL, USA).

We assessed the effect of drainage treatments on yield and GHG fluxes from rice systems by selecting only the subset of experiments that included side-by-side comparisons between continuously flooded and drained fields and repeating all meta-analyses on the reduced data set ($n = 7$). We used the natural log (lnR) of the response ratio as our effect size (Hedges *et al.*, 1999):

$$\ln R = \ln(D/F) \quad (3)$$

where D is the mean value of yield or GHG fluxes in the drained treatment and F is the mean value in the flooded treatment. To ease interpretation, the results for the analyses on lnR were back-transformed and reported as percentage change under drained conditions relative to flooded conditions ($[R-1]*100$). Treatment effects were considered significant if the 95% CI did not overlap with zero.

Results

Yields for each crop across sites averaged 6.1, 4.8, and 8.0 Mg ha⁻¹, for rice, wheat, and maize, respectively. These values were higher than the reported global averages for these crops, but showed the same overall trend

Table 2 Studies used in the meta-analysis to determine the GWP associated with greenhouse gas emissions for the major cereals

Crop	Reference	Location	CH ₄	N ₂ O	seasons of data	o*	N rate (kg ha ⁻¹)	Grain yield (Mg ha ⁻¹)	GWP range (kg CO ₂ eq season ⁻¹)	Water mgmt. †	Experimental treatments
Rice	Abao <i>et al.</i> , 2000	Los Baños, Philippines	Y	Y	2	6	30–90	5.1–6.0	75–915	CF	Residue mgmt./slow release urea
	Bhatia <i>et al.</i> , 2005	New Delhi, India	Y	Y	1	6	0–120	3.4–6.4	1027–1679	Dr	N rate/manure and residue mgmt.
	Bronson <i>et al.</i> , 1997	Los Baños, Philippines	Y	Y	3	12	80–200	4.7–6.6	174–9812	CF/Dr	N source/residue mgmt./water mgmt.
	Cai <i>et al.</i> , 1997	Jiangsu, China	Y	Y	1	5	0–300	5.2–7.3	1360–2289	Dr	N rate/N source
	Datta <i>et al.</i> , 2009	Cuttack, India	Y	Y	1	4	40	3.0–4.5	1438–2632	CF	Variety/fish
	Ghosh <i>et al.</i> , 2003	New Delhi, India	Y	Y	1	7	0–120	3.8–5.9	624–983	CF	N source/ENF*
	Kreye <i>et al.</i> , 2007	Beijing, China	Y	Y	2	6	225	3.3–8.5	177–1762	Dr	Water mgmt./plastic ground covers/mulches
	Li <i>et al.</i> , 2009	Jiangsu, China	Y	Y	1	4	300	7.9–9.4	1691–2922	Dr	ENF
	Ma <i>et al.</i> , 2007	Dapu, Jiangsu, China	Y	Y	3	18/12	0–270	5.6–8.1	685–20891	Dr	N rate/residue mgmt.
	Ma <i>et al.</i> , 2009	Dapu, Jiangsu, China	Y	Y	2	6/4	270	6.6–7.5	1455–20734	Dr	Residue mgmt.
Wheat	Ma <i>et al.</i> , 2009	Xingxiang, Jiangsu, China	Y	Y	1	5	300	5.3–6.7	2254–7393	Dr	Residue mgmt.
	Malla <i>et al.</i> , 2005	New Delhi, India	Y	Y	1	7	120–132	5.9–6.9	838–1097	Dr	ENF
	Qin <i>et al.</i> , 2010	Jiangsu, China	Y	Y	1	6	100	4.4–6.5	1430–3252	CF/Dr	Water mgmt./organic vs. mineral N
	Shang <i>et al.</i> , 2011	Hunan, China	Y	Y	6	36/12	0–204	2.2–8.8	3487–22237	CF/Dr	N,P,K mgmt./green manure
	Towprayoon <i>et al.</i> , 2005	Samutsakorn, Thailand	Y	Y	1	4	119	3.9–4.4	4035–6201	CF/Dr	Water mgmt.
	Zhang <i>et al.</i> , 2010	Jiangsu, China	Y	Y	1	6	0–300	8.6–10.2	2446–4032	Dr	Biochar/N rate
	Zou <i>et al.</i> , 2005	Jiangsu, China	Y	Y	3	10	150–479	6.6–7.9 ^s	1476–5711	CF/Dr	Water mgmt./N rate/residue mgmt.
	Abao <i>et al.</i> , 2000	Los Baños, Philippines	Y	Y	2	2	60–90	1.1–1.4	287–302	RF	Residue mgmt.
	Aulakh <i>et al.</i> , 2001a	Ludhiana, India	N	Y	1	4	0–120	1.9–5.1 [†]	1077–1264	IR	Residue mgmt.
	Bhatia <i>et al.</i> , 2005	New Delhi, India	N	Y	1	6	0–120	2.5–4.6	168–432	IR	N rate/manure and residue mgmt.
Wheat	Bhatia <i>et al.</i> , 2010	New Delhi, India	N	Y	1	10	0–120	2.3–5.6	147–409	IR	ENF/tillage
	Chen <i>et al.</i> , 2008	Jiangsu, China	N	Y	3	18	0–300	1.4–8.0 ^{ll}	1217–4349	RF	N rate/tillage/crop rotation/residue mgmt.
	Chirinda <i>et al.</i> , 2010a	Flakkebjerg, Denmark	N	Y	1	4	101–170	2.8–7.6	332–641	RF	Organic vs. conventional
	Chirinda <i>et al.</i> , 2010a,b	Foulum, Denmark	N	Y	1	5	0–165	2.8–9.5	295–431	IR	Organic vs. conventional
	Grandy <i>et al.</i> , 2006	Michigan, USA	N	Y	2	4	56–71	3.0–4.4	214–2478	RF	Tillage
	Kaiser <i>et al.</i> , 1998	Braunschweig, Germany	N	Y	3	9	0–210	6.2–10.8	515–1639	RF	N rate
	Kessavalou <i>et al.</i> , 1998	Colorado, USA	Y	Y	1	3	0	2.4–3.1	32–35	RF	Tillage
	Liu <i>et al.</i> , 2011	Shanxi, China	N	Y	1	2	180	5.6–6.1	702–749	IR	Residue mgmt.

Table 2 (continued)

Crop	Reference	Location	CH ₄	N ₂ O	seasons of data	o*	N rate (kg ha ⁻¹)	Grain yield (Mg ha ⁻¹)	GWP range (kg CO ₂ eq season ⁻¹)	Water mgmt. †	Experimental treatments
	Ma <i>et al.</i> , 2010b	Jiangsu, China	N	Y	2	8/4	138	5.0–7.2	740–1522	RF	Residue mgmt.
	Majumdar <i>et al.</i> , 2002	New Delhi, India	N	Y	1	6	0–120	2.0–4.0	351–670	IR	ENF
	Malhi & Lemke, 2007	Saskatchewan, Canada	N	Y	1	2	0–120	1.9–3.3	47–464	RF	N rate
	Malla <i>et al.</i> , 2005	New Delhi, India	N	Y	1	7	120–132	4.7–5.3	220–309	IR	ENF
	Pathak <i>et al.</i> , 2002	New Delhi, India	N	Y	1	8	0–120	3.4–5.3	144–373	IR	Water mgmt./ENF/manure
	Pathak <i>et al.</i> , 2006	New Delhi, India	N	Y	1	6	120–180	4.2–4.7	175–223	IR	Residue mgmt.
	Ruser <i>et al.</i> , 2001	Scheyern, Germany	N	Y	2	2	90–180	5.3–7.8	1262–1703	RF	N rate
	Wagner-Riddle <i>et al.</i> , 2007	Ontario, Canada	N	Y	1	2	60–90	6.8–7.6**	318–387	RF	N rate
	Webb <i>et al.</i> , 2004	Top Kingston, UK	N	Y	2	2	180–190	4.7–7.5	328–562	RF	Cropping system/annual variation
	Webb <i>et al.</i> , 2004	Welbeck, UK	N	Y	2	2	180–190	4.2–6.0	468–515	RF	Cropping system/annual variation
	Webb <i>et al.</i> , 2004	Propagation, UK	N	Y	3	3	150–190	9.0–10.2	515–562	RF	Cropping system/annual variation
	Webb <i>et al.</i> , 2004	Shepard's Gate, UK	N	Y	3	3	160–230	8.4–10.0	515–843	RF	Cropping system/annual variation
	Wei <i>et al.</i> , 2010	Shanxi, China	N	Y	2	10/5	0–120	1.3–5.4	546–2407	RF	Fertilizer mgmt./manure
	Weiske <i>et al.</i> , 2001	Giessen, Germany	Y	Y	1	3	180	6.4–6.7	77–165	RF	ENF
Maize	Adviento-Borbe <i>et al.</i> , 2007	Nebraska, USA	Y	Y	2	10	140–310	12.0–17.5	522–2245	IR	N rate/crop rotation
	Adviento-Borbe <i>et al.</i> , 2010	Pennsylvania, USA	N	Y	2	8/4	90–225	10.6–15.1	468–4167	RF	N rate/crop rotation/manure
	Almaraz <i>et al.</i> , 2009	Quebec, Canada	N	Y	1	4	0–180	2.8–7.9	1072–2580	RF	N rate/tillage
	Guo <i>et al.</i> , 2009	Hubei, China	N	Y	1	6	0	1.8–3.6	136–253	RF	Alley cropping/cropping system
	Halvorson <i>et al.</i> , 2010	Colorado, USA	Y	Y	2	14/7	0–246	7.9–14.3	71–406	IR	ENF
	Khalil <i>et al.</i> , 2002	University Putra, Malaysia	N	Y	1	3	150–243	1.0–1.4	351–1241	RF	Residue mgmt./manure
	Liu <i>et al.</i> , 2011	Shanxi, China	N	Y	1	2	210	7.5–7.6	890–1405	IR	Residue mgmt.
	McSwiney & Robertson, 2005	Michigan, USA	N	Y	3	27/9	0–291	3.2–9.4	453–4721	IR/RF	N rate
	Mosier <i>et al.</i> , 2006	Colorado, USA	Y	Y	3	22/11	0–224	4.2–12.7	93–1869	IR	N rate/tillage/crop rotation
	Parkin & Hatfield, 2010	Iowa, USA (1)	N	Y	1	2	125	10.7–11.8	2781–3291	RF	ENF
	Parkin & Hatfield, 2010	Iowa, USA (2)	N	Y	1	2	168	12.4–12.7	2463–2720	RF	ENF

Table 2 (continued)

Crop	Reference	Location	CH ₄	N ₂ O	seasons of data	o*	N rate (kg ha ⁻¹)	Grain yield (Mg ha ⁻¹)	GWP range (kg CO ₂ eq season ⁻¹)	Water mgmt.†	Experimental treatments
	Parkin & Kaspar, 2006	Iowa, USA	N	Y	2	3	215	9.7–10.0	4490–5389	RF	Tillage/residue mgmt.
	Phillips <i>et al.</i> , 2009	North Dakota, USA	Y	Y	1	2	70	2.5–2.7	158–203	RF	N timing
	Qian <i>et al.</i> , 1997	Nebraska, USA	N	Y	2	2	153–234	10.3–10.4	712–1072	IR	High nitrate water
	Ruser <i>et al.</i> , 2001	Scheyern, Germany	N	Y	2	2	65–130	8.0–9.5	585–998	RF	N rate
	Sehy <i>et al.</i> , 2003	Scheyern, Germany	N	Y	1	4	125–175	8.9–10.5	1170–3558	RF	Site specific fertilization
	Venterea <i>et al.</i> , 2010	Minnesota, USA	N	Y	3	12/4	146	6.6–10.5	281–1592	RF	N source/crop rotation
	Wagner-Riddle <i>et al.</i> , 2007	Ontario, Canada	N	Y	2	4	50–150	3.9–9.0**	468–824	RF	N source and mgmt.
	Weiske <i>et al.</i> , 2001	Giessen, Germany	Y	Y	1	3	160	7.7–8.0	85–159	RF	ENF
	Yan <i>et al.</i> , 2001	Tsukuba, Japan	N	Y	1	4	0–250	2.0–2.5	59–255	RF	N mgmt./ENF

*Indicates number of observations. A number following a '/' indicates the number of observations used in analysis after averaging similar treatments across years.

†Water management: Continuous flood (CF); rice with any drainage or aerobic period (Dr); irrigated (IR); rainfed (RF).

‡ENF, enhanced-efficiency N fertilizer (includes nitrification inhibitors, urease inhibitors, slow release fertilizers).

§Yield data: personal communication (R. Sass).

¶Aulakh *et al.* (2001b).

||Calculated by equation (Chen *et al.*, 2008; Fig 2a).

**Jayasundara *et al.* (2007).

Y, yes; N, no.

with maize yields being the highest and wheat the lowest (Table 1). Yields ranged between 2.2 and 10.2 Mg ha⁻¹ for rice, 1.1 and 10.8 Mg ha⁻¹ for wheat, and 1.0 and 17.5 Mg ha⁻¹ for maize (Table 2). The average N rate applied was 172, 115, and 152 kg N ha⁻¹ for rice, wheat, and maize, respectively (data not shown). The relatively high average N rate for rice is in part due to 10 of the 17 study sites being from China. Fertilizer N rates in China are known to be excessive (Ju *et al.*, 2009) and in our database, China was the only country that had N rates in excess of 200 kg N ha⁻¹ in rice systems (Table 2).

GHG fluxes, GWP and yield-scaled GWP for each crop

The average GWP of CH₄ and N₂O emissions was highest for rice systems (3757 kg CO₂ eq ha⁻¹ season⁻¹), which was 5.7 times higher than for wheat (662 kg CO₂ eq ha⁻¹ season⁻¹) and 2.7 times higher than for maize (1399 kg CO₂ eq ha⁻¹ season⁻¹) (Fig. 1). The GWP coefficient of variation across all observations for each crop was high, ranging from 0.98 to 1.13, with values across crops being similar. The range in GWP across all observations for each crop was 75–22 237 kg CO₂ eq ha⁻¹ season⁻¹ for rice, 32–4349 kg CO₂ eq ha⁻¹ season⁻¹ for wheat, and 59–5389 kg CO₂ eq ha⁻¹ season⁻¹ for maize (Table 2).

The GWP of CH₄ and N₂O emissions from rice systems was largely determined by CH₄ emissions (100 kg CH₄-C ha⁻¹ season⁻¹) that accounted for 89% of the GWP. Three rice studies (Ma *et al.*, 2007, 2009; Shang *et al.*, 2011), all from China, reported extremely high CH₄ emissions in excess of 20 000 kg CO₂ eq ha⁻¹ sea-



Fig. 1 Results of a meta-analysis on net seasonal soil fluxes of CH₄ and N₂O from three cereal crops. For N₂O fluxes, the results for rice, wheat and maize were based on 116, 122, and 88 observations, respectively. For CH₄ fluxes, the results were based on 116, 33, and 8 observations. Observations were weighted by sampling frequency and replication. All error bars represent 95% confidence intervals.

son⁻¹, whereas none of the other studies reported values in excess of 10 000 kg CO₂ eq ha⁻¹ season⁻¹ (Table 2). In the few studies that reported CH₄ emissions in wheat and maize, CH₄ emissions represented <2% of GWP, and on average these systems were a minor sink for CH₄ (-0.3 kg CH₄-C ha⁻¹ season⁻¹). All cropping systems were net emitters of N₂O, with rice emitting the least (0.88 kg N₂O-N ha⁻¹ season⁻¹) followed by wheat (1.44 kg N₂O-N ha⁻¹ season⁻¹) and maize (3.01 kg N₂O-N ha⁻¹ season⁻¹).

The yield-scaled GWP was significantly higher for rice (657 kg CO₂ eq Mg⁻¹) than for wheat and maize (Fig. 2). Despite higher GWP of emissions in maize, the yield-scaled GWP was similar for both wheat (166 kg CO₂ eq Mg⁻¹) and maize (185 kg CO₂ eq Mg⁻¹) due to lower wheat yields (Table 1).

Methane, nitrous oxide and GWP in relation to N input

There was no effect of N rate on CH₄ emissions in rice (Fig. 3b); however, there was a significant correlation between N input and N₂O emissions for all crops (Fig 3a,c,d), although rice N₂O emissions were roughly 60% of that for wheat and maize. On average, 0.68%, 1.21%, and 1.06% of N applied was emitted as N₂O in rice, wheat, and maize, respectively. Yields for all crops increased with increasing N rate, although to a lesser extent in rice systems (Table 3). The yield-scaled GWP was not affected by N rate in rice systems; however, in maize and wheat systems, it tended to be similar for low and medium N rates, but increased at the highest N rates (although this trend was not always significant).




Fig. 2 Results of a meta-analysis on yield-scaled GWP for three cereal crops. The results for rice, wheat, and maize were based on 116, 122, and 88 observations, respectively. CH₄ flux data were available for only a few of the studies on wheat and maize. As CH₄ fluxes were negligible for these two crops, they were not included in GWP calculations. Observations were weighted by sampling frequency and replication. All error bars represent 95% confidence intervals.




Fig. 3 Net seasonal soil fluxes of N_2O and CH_4 vs. N application rate for rice (a,b), and net seasonal soil fluxes of N_2O vs. N application for wheat (c) and maize (d).

Discussion

Meta-analysis results compared to previous estimates

To our knowledge, no published estimates of global N_2O emissions exist for individual crops, including the major cereals rice, wheat, and maize. However, a number of studies have attempted to estimate global CH_4 emissions from rice systems. Recently, Yan *et al.* (2009) reported that previous estimates have ranged from 25.6 Tg $\text{CH}_4 \text{ yr}^{-1}$ (Yan *et al.*, 2009) to 120 Tg $\text{CH}_4 \text{ yr}^{-1}$ (Holzapfel-Pschorn & Seiler, 1986), with the IPCC using an estimate of 60 Tg $\text{CH}_4 \text{ yr}^{-1}$ (IPCC, 1995). For comparison, we used the mean CH_4 emission value from our analysis of 134 kg $\text{CH}_4 \text{ ha}^{-1} \text{ season}^{-1}$ (100 kg $\text{CH}_4\text{-C ha}^{-1} \text{ season}^{-1}$) and multiplied by the total harvested rice area in (Table 1) to estimate global CH_4 emissions of 21.6 Tg $\text{CH}_4 \text{ yr}^{-1}$, or roughly one-third of current IPCC estimates. Our value was similar to that of Yan *et al.* (2009), where they estimated that in most Asian rice growing countries, the percent area with one or more drainage events ranged from 57% to 80%. This estimate is roughly in line with our analysis in which 76% of the observations used some form of mid-season drain and/or intermittent irrigation.

For wheat and maize systems, our results indicate that 1.21% and 1.06%, respectively, of applied N is emitted as N_2O (Fig. 3c,d); slightly higher than the

suggested 1% used by the IPCC (IPCC, 2006). In flooded rice systems, N_2O emissions resulting from applied N are expected to be lower than for upland crops, and we found that 0.68% of applied N was emitted as N_2O . Based on a review, Akiyama *et al.* (2005) reported N_2O fertilizer emission factors of 0.22% for continuously flooded rice systems and 0.37% for intermittently flooded rice systems. The IPCC guidelines estimate that on average 0.3% of N fertilizer applied to rice paddies is emitted as N_2O (IPCC, 2006). The higher values reported from our analysis may be due to the large number of studies from China, where fertilizer N rates for rice systems are generally high (Ju *et al.*, 2009). Such rates were evident in the studies used for this analysis (Table 2). One reason for lower N_2O emissions in rice systems compared with maize and wheat systems is because rice soils are often submerged, so a large portion of the N_2O that is produced is further reduced to N_2 (Firestone & Davidson, 1989; Hou *et al.*, 2000; Aulakh *et al.*, 2001a). Most N_2O emissions from rice systems occur during drainage events when NH_4^+ is converted to NO_3^- , which then becomes susceptible to denitrification (Yao *et al.*, 2010). Indeed, in our analysis, the average N_2O emissions from continuously flooded rice systems was only 0.19 kg $\text{N}_2\text{O-N ha}^{-1} \text{ season}^{-1}$ (data not shown) compared with the overall rice average that was 0.88 kg $\text{N}_2\text{O-N ha}^{-1} \text{ season}^{-1}$ (Fig. 1).

Table 3 Yield, GWP for net seasonal GHG fluxes, and yield-scaled GWP (GWP/yield) for three different cereal crops, as affected by N fertilization rate. As CH₄ fluxes were negligible for wheat and maize, GWP for these crops were calculated from net seasonal N₂O fluxes alone

	N rate kg N ha ⁻¹ yr ⁻¹	Mean N rate	Yield Mg ha ⁻¹ season ⁻¹	95% CI season ⁻¹	GWP (CH ₄) kg CO ₂ eq ha ⁻¹ season ⁻¹	95% CI season ⁻¹	GWP (N ₂ O) kg CO ₂ eq ha ⁻¹ season ⁻¹	95% CI season ⁻¹	GWP/yield kg CO ₂ eq Mg ⁻¹	95% CI
Rice	0-50	13.3	5.14	4.29-6.26	3124	1943-5191	151	93-209	678	428-1069
	50-125	109.9	5.24	4.96-5.50	3117	2173-4321	173	124-240	672	479-910
	125-200	176.8	6.45	6.18-6.74	3944	2047-6423	120	75-177	630	344-982
	>200	282.2	7.05	6.53-7.53	3496	2166-5290	802	585-1049	643	450-889
Wheat	0-50	0	2.53	2.19-2.97	-	-	340	173-638	165	78-316
	50-125	107.3	4.48	4.07-4.96	-	-	573	448-743	162	113-237
	125-200	170.4	6.11	5.74-6.50	-	-	887	650-1191	159	115-220
	>200	252.5	6.27	4.44-9.16	-	-	1808	945-3109	369	152-602
Maize	0-50	1.6	4.40	3.21-6.16	-	-	441	147-890	131	39-304
	50-125	90.7	4.13	2.91-7.54	-	-	527	280-1203	102	85-140
	125-200	150.5	9.31	7.40-10.68	-	-	1622	937-2182	183	127-232
	>200	223.7	9.16	7.59-10.80	-	-	1827	1087-2731	246	148-374

*Indicates number of observations.

Differences in GWP between crops

Results from our meta-analysis show that seasonal GWP ha⁻¹ is the highest in rice systems, followed by maize and then wheat. Differences in N₂O emissions between maize and wheat are most likely due to differences in N input, which averaged 115 kg ha⁻¹ in wheat systems compared with 152 kg ha⁻¹ in maize systems. The GWP of GHG emissions for wheat and maize was almost entirely driven by N₂O emissions that are related to fertilizer N input. Although an analysis of the data across all studies support this finding (Fig. 3c,d), data from individual studies show a much stronger correlation between N rate and N₂O emission (e.g. McSwiney & Robertson, 2005; Chen *et al.*, 2008). It is possible that the GWP of N₂O emissions for wheat may be over-estimated relative to the other crops in this analysis, as most of the studies were conducted on winter wheat and roughly a third of these winter wheat studies were conducted in temperate climates. As our estimate of GWP covers the period from planting to harvest, winter emissions are included. Some have reported that winter N₂O emissions can account for up to half of annual emissions (Kaiser *et al.*, 1998).

The differences in the GWP of emissions between wheat and maize are small relative to the difference between rice and the other two crops. Rice systems emit N₂O, which were related to N input, but N₂O emissions for rice were significantly lower than for either wheat or maize and contributed only 11% to total GWP of rice emissions. The main difference between rice and wheat or maize systems was the high CH₄ emissions for rice (Fig. 1). Rice systems are fundamentally different from wheat and maize systems for several reasons. First, rice is typically grown in flooded soils creating anaerobic conditions leading to methanogenesis, which is a strictly anaerobic microbial decomposition process of organic material. Second, rice plants are important conduits of GHGs (both CH₄ and N₂O) from the soil to the atmosphere (Yu *et al.*, 1997). Methane transport through the plant is the dominant form of methane release into the atmosphere and can account for up to 90% of total emissions (Holzapfel-Pschorn & Seiler, 1986; Butterbach-Bahl *et al.*, 1997). Rice systems are relatively efficient at oxidizing CH₄, with an estimated 58% (Sass *et al.*, 1990) to 80% (Holzapfel-Pschorn *et al.*, 1986; Conrad & Rothfuss, 1991) of the CH₄ produced in the soil being oxidized by methanotrophs and not emitted to the atmosphere. Interestingly, the high GWP of CH₄ and N₂O emissions for rice is despite the fact that 14 of the 17 studies applied some form of drainage (Table 2). As will be discussed later, field drainage is often recommended as a mitigation strategy to reduce CH₄ emissions in rice systems.

Table 4 Results of a meta-analysis on yield, GWP for net seasonal GHG fluxes, and yield-scaled GWP for three different cereal crops. For each experimental site in our dataset, we only considered observations from plots with the lowest yield-scaled GWP. As CH₄ fluxes were negligible for wheat and maize, GWP for these crops were calculated from net seasonal N₂O fluxes alone

Crop	o*	N rate kg N ha ⁻¹ season ⁻¹	Grain yield Mg ha ⁻¹ season ⁻¹	95% CI	% of maximum yield†	GWP (CH ₄) kg CO ₂ eq ha ⁻¹ season ⁻¹	95% CI	GWP (N ₂ O)	95% CI	Yield- scaled GWP	95% CI
Rice	17	193	6.64	5.94–7.40	94	1301	797–2011	313	202–444	279	169–439
Wheat	25	101	5.04	4.20–5.87	91	–	–	595	280–1168	102	61–169
Maize	20	119	7.60	5.34–9.47	90	–	–	1115	445–1819	140	82–209

*Indicates number of observations.

†Percent of maximum yield was calculated as yield with the lowest yield-scaled GWP/maximum yield obtained per site.

Yield-scaled GWP: crops and N input

Focusing on reducing GWP on an area basis, as is commonly done, can be counterproductive, because it can lead to low yielding management practices. In contrast, yield-scaled GWP is an integrated metric that addresses the dual goals of environmental protection and food security. Yield-scaled GWP was significantly higher for rice than either wheat or maize, which were comparable. Therefore, these results lead us to reject our first hypothesis (that yield-scaled GWP for the major cereals are similar). To our knowledge, no reviews have been conducted that compare crops in this manner. However, Pathak *et al.* (2010) recently evaluated the C footprint of Indian food items and reported that yield-scaled GWP of rice was approximately 10 times that of wheat, whereas in our meta-analysis yield-scaled GWP of rice was about four times higher than wheat (Fig. 2). Two of the studies in our analysis (Bhatia *et al.*, 2005; Malla *et al.*, 2005) were from Indian rice–wheat systems, where GHG emissions were measured in both rice and wheat crops. They found that, on average, the yield-scaled GWP of rice was three times higher than wheat (210 kg CO₂ eq Mg⁻¹ and 73 kg CO₂ eq Mg⁻¹, respectively, data not shown), suggesting the GWP estimates in Pathak *et al.* (2010) are unexpectedly high and not supported by our findings.

Our data show a trend that yield-scaled GWP for maize and wheat remain relatively constant at low to moderate N rates, but increases when N fertilizer rates are high (Table 3). Other studies have also reported that N rates in excess of crop N demand can lead to large N₂O losses (McSwiney & Robertson, 2005; Ma *et al.*, 2010a) and higher yield-scaled GWP (van Groenigen *et al.*, 2010). In agreement, Hoben *et al.* (2011) recently suggested that maize farmers should be encouraged to apply N at a rate sufficient for maximum economic returns, and that applying N at higher rates to achieve maximum agronomic returns leads to only marginal yield increases, but much higher N₂O emissions.

In rice, the GWP of emissions is primarily driven by CH₄ and not N₂O. Previous work has shown that increasing the N rate through the use of urea has the potential to both increase (Lindau *et al.*, 1991; Corton *et al.*, 2000) and decrease (Zou *et al.*, 2005; Xie *et al.*, 2010) CH₄ emissions. Our study shows no correlation between N rate and CH₄ emissions (Fig. 3b) and higher N₂O emissions at excessive N rates (Table 3), leading to a recommendation of achieving maximum economic returns to reduce yield-scaled GWP, similar to wheat and maize.

Table 5 Summary of the results of the meta-analysis on the effect of drainage on GWP for net seasonal GHG fluxes, yield and yield-scaled GWP (GWP/yield) for rice, using the response metric $\ln R$ (see Methods). The analysis is based on seven side-by-side comparisons between drained and undrained treatments

	Average effect (%)	95% CI
GWP	-33.6	-42.9 to -23.7
Yield	0.95	-4.69 to 7.63
GWP/yield	-34.3	-45.6 to -23.4
N ₂ O	445	121 to 1502
CH ₄	-49.5	-58.8 to -26.3

Opportunities to reduce GWP and yield-scaled GWP

It is not within the scope of this analysis to identify management practices that result in the lowest yield-scaled GWP for these crops; however, our data set provides insight on the potential to reduce yield-scaled GWP. In rice systems, the average lowest yield-scaled GWP was 279 kg CO₂ eq Mg⁻¹, whereas in wheat and maize, these values were 102 and 140 kg CO₂ eq Mg⁻¹, respectively (Table 4). Thus, under this 'best case' scenario, the yield-scaled GWP for rice is about twice as high as for wheat or maize, which is an improvement on the overall average where the yield-scaled GWP was about 3.75 times higher for rice than wheat or maize (Fig. 2).

The yield penalty (i.e. the yield reduction relative to the highest yield for any given site) to achieve the lowest yield-scaled GWP was 6%, 9% and 10% for rice, wheat, and maize, respectively (Table 4). Such results are encouraging when we recognize the need for high yields to achieve food security (Tilman *et al.*, 2002; Lobell *et al.*, 2009; Burney *et al.*, 2010). These data support that of others (Robertson & Vitousek, 2009; Hoben *et al.*, 2011) who have found that substantial N₂O emissions can be avoided by improved N management strategies that better match crop N demand and economic returns, as opposed to applying N for maximum agronomic yields (at N rates which tend to be higher than to achieve maximum economic returns). Such strategies provide a potential win-win for growers and the environment.

In rice systems, draining a field at some point during the growing season is commonly recommended as a means of reducing CH₄ emissions (e.g., Yagi *et al.*, 1997; Wassmann *et al.*, 2000). However, draining can also increase N₂O emissions (Hou *et al.*, 2000) which may outweigh benefits to reduction of overall GWP, as N₂O is a much more potent GHG. However, most studies report that the GWP (accounting for both CH₄ and N₂O) in drained rice fields is lower than in continuously flooded fields (e.g. Zou *et al.*, 2005). Our analysis

of studies that included side-by-side comparisons of continuously flooded and drained fields showed that drainage decreased both GWP and yield-scaled GWP by about 34%, but yields were not significantly affected (Table 5). As expected, N₂O emissions were higher and CH₄ emissions lower in drained fields. Similarly, Wassmann *et al.* (2000) reported that draining rice fields could mitigate CH₄ emissions by 7–80% depending on timing, frequency, and duration of drainage.

Limitations of our analysis

Our analysis did not include studies where crops were evaluated side-by-side in the same season (such comparisons are not feasible as these crops have different climatic requirements for optimal production); therefore, direct comparisons among crops were not possible. However, our analysis included a number of studies where one crop followed another in a rotation, meaning the crops were grown on the same soil. An example of this is the rice–wheat rotation (Bhatia *et al.*, 2005; Malla *et al.*, 2005). In these two studies, the GWP of CH₄ and N₂O emissions for rice systems was three times higher than wheat, roughly similar to our meta-analytic approach, where it was four times that of wheat.

Most of the studies in our data set were conducted at experimental sites under the management of researchers. Thus, the management and environment may not always be representative of actual production fields. Also, in several of these studies, treatments were evaluated that are not being widely used by farmers. Examples of this include the use of enhanced-efficiency N fertilizer (ENF) (Table 2), plastic mulches (Kreye *et al.*, 2007), and alley cropping (Guo *et al.*, 2009). As most of these treatments were aimed at reducing GWP, it is possible that our overall estimates may be biased toward low GHG fluxes.

Values for annual, rather than seasonal, GHG emissions are needed for these cropping systems as fallow periods can be a large contributor to net GHG fluxes. Moreover, treatment effects like rate of N fertilizer input and its impact on residue yield and C : N ratios can last beyond the growing season. Kaiser *et al.* (1998) reported for a cereal study in Europe that approximately 50% of annual emissions occurred during the winter months (October to February) due to freezing and thawing events during this period. Unfortunately, a meta-analysis of annual GHG emissions was not possible for two reasons. First, the large majority of studies only provided GHG emission data for the growing season. Second, in many parts of the world, especially in the tropics, multiple crops are grown annually on the same piece of land. For example, rice–wheat systems

(where rice is grown during the summer and wheat in the winter) are common in both India and China and occupy 24 million ha of land (<http://www.rwc.cgiar.org>), or about 15% of the total area under rice production. Thus, annual comparisons of GHG emissions from different crops are not always possible, as different crops are grown within the same year.

The important agricultural GHGs are N₂O, CH₄, and CO₂; however, for our analysis, most of the wheat and maize studies did not include measurements of CH₄ and CO₂. On a global scale, soil CO₂ fluxes are largely offset by high rates of net primary productivity and atmospheric CO₂ fixation by crop plants, and thus contribute <1% to the GWP of agriculture (Smith *et al.*, 2007). It is generally agreed that the net balance between C respiration and fixation in a cropping system is reflected by changes in soil organic carbon over time (West & Post, 2002; Stewart *et al.*, 2007). However, within a typical study period of 1–2 years (as is the case with most GHG studies), differences in soil organic carbon are difficult to detect as the magnitude of change is small and there is a large degree of spatial variability (Post *et al.*, 2001; Conant *et al.*, 2011). Long-term studies, however, have shown that soil C sequestration or loss can significantly influence net GWP (Robertson *et al.*, 2000; Six *et al.*, 2004) and thus cannot be neglected. In relation to our objectives, reports out of China have found that rice systems have a higher potential for C sequestration than crops grown under aerobic conditions (Pan *et al.*, 2010; Wu, 2011).

To be included in our dataset, wheat and maize studies did not have to report CH₄ emissions. Only a few studies measured CH₄ emissions and they were minor compared with N₂O emissions. Methane is produced in agricultural soils exclusively by a group of bacteria known as methanogens, via the anaerobic process of methanogenesis that occurs at redox potentials less than –150 mV (Masscheleyn *et al.*, 1993; Mosier *et al.*, 2004). Methanogenesis is a strictly anoxic process and therefore upland cropping systems are not a direct sources of CH₄ under normal conditions (Bronson & Mosier, 1993; Robertson & Grace, 2004), although some studies have reported CH₄ emissions from upland cropping systems having values as high as 0.6 kg CH₄-C ha⁻¹ season⁻¹ (Abao *et al.*, 2000; Mosier *et al.*, 2006; Halvorson *et al.*, 2010). In contrast, other studies found that the net flux of CH₄ in upland soils can be negative, if it is largely a function of CH₄ oxidation, where CH₄ is oxidized to CO₂ by methanotrophic bacteria under oxic conditions. (Robertson *et al.*, 2000; Hütsch, 2001; Robertson & Grace, 2004; Ellert & Janzen, 2008). Although the maximum CH₄ oxidation rate in wheat and

maize systems was determined to be 2.19 kg CH₄-C ha⁻¹ season⁻¹ for maize (Adviento-Borbe *et al.*, 2007), overall, the effect of CH₄ oxidation was minor and represented <2% of GWP.

Finally, the intent of this analysis was not to identify the most sustainable crop with respect to land use decisions and climate change, but rather to compare the GWP of CH₄ and N₂O emissions of major cereals based on field GHG measurements and broadly determine if the lowest yield-scaled GWP for each crop was achieved at near optimal yields. Our results support the conclusion that lower yield-scaled GWP values can be achieved at near optimal yields. Addressing the issue of environmental sustainability for each crop is important and new approaches and tools are being developed to more comprehensively assess the impacts of agricultural management decisions using multiple indicators (Steffan-Dewenter *et al.*, 2007; DeFries & Rosenzweig, 2010); however, this was not the objective of this study. Emissions of GHG from soil are only one of many factors that need to be considered when determining the net C footprint of agricultural systems and evaluating potential mitigation practices to be implemented (Smith *et al.*, 2007). Certainly, one other consideration is the C sequestration potential of each system, as discussed above. Another important consideration is the percentage of these cereals that are consumed directly by humans. Rice is the staple crop for the largest number of people on earth, and human consumption accounts for 85% of total production of rice. In contrast, 72% of wheat and only 19% of maize is directly consumed by humans (Maclean *et al.*, 2002). A larger proportion of these crops are used for other purposes such as livestock feed and biofuels, which in turn have associated GHG emissions and energy requirements that would need to be factored into a complete life cycle analysis to more fully understand and account for environmental impacts among cereal crops (Hill *et al.*, 2006; Garnett, 2009). Other considerations that would need to be addressed, but are much more difficult to quantify, are the ecosystem services these crops provide and the roles of these crops in various cultures (Steffan-Dewenter *et al.*, 2007; Sachs *et al.*, 2010). For example, in California, 230 wildlife species and an estimated 10–12 million water birds use rice fields annually (Sterling & Buttner, 2009). It is increasingly recognized that integrated metrics involving ecosystem services, GWP estimates, and other important aspects of agriculture (e.g. human health, food security, economic prosperity, and sociocultural well-being) need to be developed to begin to monitor and compare the impacts of agricultural practices on a global scale (Sachs *et al.*, 2010).

Need for comparable GHG emission estimates

When conducting this meta-analysis, it was apparent that most studies only discussed results for GHG emissions relative to other treatments in their study rather than to other comparable studies. One possible reason for this is that, to our knowledge, there are no established or well-accepted values for comparison of GWP results between studies. In this meta-analysis, we made an attempt to provide average values for GWP, calculated per ha and per ton of grain produced (yield-scaled). A second problem is that many different units are used to express GWP that makes straightforward comparisons with other studies difficult. When compiling this data set, a combination of the following units was used: CH₄ (or CH₄-C), N₂O (or N₂O-N) µg, mg, kg, h⁻¹ day⁻¹, season⁻¹, and year⁻¹. To allow for better comparisons to be made between values in the future, we propose for field studies that GHG be reported as kg CH₄-C (or N₂O-N) ha⁻¹ season⁻¹ (or year⁻¹, depending on study) and that yield-scaled GWP be reported as kg CO₂ eq Mg⁻¹.

Interestingly, the primary limitation to compiling a larger data set for this analysis was the absence of yield data in studies where GHG were measured. As there are concerns about global food security and how mitigation strategies may affect yield, it will become increasingly important to link the intensity of GHG emissions with the primary function of these systems (i.e. producing food). Therefore, when GHG studies are conducted in cropping systems, yield data should be routinely provided to facilitate further and more in-depth analysis on the relationship between mitigation practices, GHG emissions, and yield.

Conclusions

We conducted a comprehensive meta-analysis of GHG emissions in major cereal cropping systems and related GWP with grain yield. Our results show that the GWP of CH₄ and N₂O emissions from rice was significantly higher than for wheat or maize when expressed on an area basis. Likewise, yield-scaled GWP remained about 3.75 times higher for rice compared with wheat and maize. The higher GWP of emissions from rice were largely driven by CH₄ emissions, which were unaffected by fertilizer N input. However, of the three cereals evaluated, rice systems showed the greatest potential for reduction in yield-scaled GWP. The lowest yield-scaled GWP values for each crop were achieved at near optimal yields, highlighting the potential for sustainable intensification of these cereals that will be necessary to meet the increased demand brought on by increasing population. There was also evidence that relatively small

increases in yield, in particular, for wheat and maize through increased N fertilization, led to disproportionately higher yield-scaled GWP. This finding indicates that better N management practices would allow for a reduction in yield-scaled GWP without significant yield penalties. Our analysis showed that mitigation practices to reduce yield-scaled GWP for rice should be mainly focused on the reduction of CH₄ emissions, even though these mitigation practices will probably lead to higher N₂O emissions. Importantly, we were not able to evaluate the C sequestration potential of these systems in our analysis and some studies have shown that rice systems have a greater potential to sequester C than upland cropping systems. Finally, GWP is only one of many factors that need to be considered when evaluating the sustainability of different cereal crops. For example, more rice is directly consumed by humans than either wheat or maize, where additional GHG emissions and energy costs may be associated with subsequent conversion of these crops into feed or fuel. In the context of climate change and agriculture, there is growing consensus that other factors including cultural significance, the provisioning of ecosystem services, food security, and human health and well-being must also be considered.

Acknowledgement

This work was generally supported by Mars Inc., and the California Rice Research Board. K.J. van Groenigen is supported by a grant from the Irish Research Council for Science, Engineering and Technology, co-funded by Marie Curie Actions under FP7. We would also like to thank the Department of Plant Sciences, UC Davis for supporting C. Pittelkow with a Graduate Student Research Assistantship Award.

References

- Abao EB, Bronson KF, Wassmann R, Singh U (2000) Simultaneous records of methane and nitrous oxide emissions in rice-based cropping systems under rainfed conditions. *Nutrient Cycling in Agroecosystems*, **58**, 131–139.
- Adviento-Borbe MAA, Haddix ML, Binder DL, Walters DT, Dobermann A (2007) Soil greenhouse gas fluxes and global warming potential in four high-yielding maize systems. *Global Change Biology*, **13**, 1972–1988.
- Adviento-Borbe MAA, Kaye JP, Bruns MA, McDaniel MD, McCoy M, Harkcom S (2010) Soil greenhouse gas and ammonia emissions in long-term maize-based cropping systems. *Soil Science Society of America Journal*, **74**, 1623–1634.
- Akiyama H, Yagi K, Yan X (2005) Direct N₂O emissions from rice paddy fields: summary of available data. *Global Biogeochemical Cycles*, **19**, GB1005.
- Almaraz JJ, Mabood F, Zhou X, Madramootoo C, Rochette P, Ma BL, Smith DL (2009) Carbon dioxide and nitrous oxide fluxes in corn ground under two tillage systems in Southwestern Quebec. *Soil Science Society of America Journal*, **73**, 113–119.
- Aulakh MS, Khera TS, Doran JW, Bronson KF (2001a) Denitrification, N₂O and CO₂ fluxes in rice-wheat cropping system as affected by crop residues, fertilizer N and legume green manure. *Biology and Fertility of Soils*, **34**, 375–389.
- Aulakh MS, Khera TS, Doran JW, Bronson KF (2001b) Managing crop residue with green manure, urea, and tillage in a rice-wheat rotation. *Soil Science Society of America Journal*, **65**, 820–827.
- Bhatia A, Pathak H, Jain N, Singh PK, Singh AK (2005) Global warming potential of manure amended soils under rice-wheat system in the Indo-Gangetic plains. *Atmospheric Environment*, **39**, 6976–6984.

- Bhatia A, Sasmal S, Jain N, Pathak H, Kumar R, Singh A (2010) Mitigating nitrous oxide emission from soil under conventional and no-tillage in wheat using nitrification inhibitors. *Agriculture, Ecosystems & Environment*, **136**, 247–253.
- Bronson KF, Mosier AR (1993) Nitrous oxide emissions and methane consumption in wheat and corn-cropped systems. In: *Agricultural Ecosystem Effects on Trace Gases and Global Climate Change, American Society of Agronomy Special Publication No. 55* (eds Harper LA, Mosier AR, Duxbury JM, Rolston DE), pp. 133–144. American Society of Agronomy, Madison, WI.
- Bronson KF, Neue HU, Singh U, Abao EB (1997) Automated chamber measurements of methane and nitrous oxide flux in a flooded rice soil: I. Residue, nitrogen, and water management. *Soil Science Society of America Journal*, **61**, 981–987.
- Burney JA, Davis SJ, Lobell DB (2010) Greenhouse gas mitigation by agricultural intensification. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 12052–12057.
- Butterbach-Bahl K, Papen H, Rennenberg H (1997) Impact of gas transport through rice cultivars on methane emission from rice paddy fields. *Plant, Cell & Environment*, **20**, 1175–1183.
- Cai Z, Xing G, Yan X, Xu H, Tsuruta H, Yagi K, Minami K (1997) Methane and nitrous oxide emissions from rice paddy fields as affected by nitrogen fertilisers and water management. *Plant and Soil*, **196**, 7–14.
- Cassman KG (1999) Ecological intensification of cereal production systems: yield potential, soil quality, and precision agriculture. *Proceedings of the National Academy of Sciences of the United States of America*, **96**, 5952–5959.
- Cassman KG, Dobermann A, Walters DT, Yang H (2003) Meeting cereal demand while protecting natural resources and improving environmental quality. *Annual Review of Environment and Resources*, **28**, 315–358.
- Chen S, Huang Y, Zou J (2008) Relationship between nitrous oxide emission and winter wheat production. *Biology and Fertility of Soils*, **44**, 985–989.
- Chirinda N, Carter MS, Albert KR, Ambus P, Olesen JE, Porter JR, Petersen SO (2010a) Emissions of nitrous oxide from arable organic and conventional cropping systems on two soil types. *Agriculture, Ecosystems & Environment*, **136**, 199–208.
- Chirinda N, Olesen JE, Porter JR, Schjønning P (2010b) Soil properties, crop production and greenhouse gas emissions from organic and inorganic fertilizer-based arable cropping systems. *Agriculture, Ecosystems & Environment*, **139**, 584–594.
- Cole CV, Duxbury J, Freney J *et al.* (1997) Global estimates of potential mitigation of greenhouse gas emissions by agriculture. *Nutrient Cycling in Agroecosystems*, **49**, 221–228.
- Conant RT, Ogle SM, Paul EA, Paustian K (2011) Measuring and monitoring soil organic carbon stocks in agricultural lands for climate mitigation. *Frontiers in Ecology and the Environment*, **9**, 169–173.
- Conrad R, Rothfuss F (1991) Methane oxidation in the soil surface layer of a flooded rice field and the effect of ammonium. *Biology and Fertility of Soils*, **12**, 28–32.
- Corton TM, Bajita JB, Grospe FS *et al.* (2000) Methane emission from irrigated and intensively managed rice fields in Central Luzon (Philippines). *Nutrient Cycling in Agroecosystems*, **58**, 37–53.
- Datta A, Nayak DR, Sinhababu DP, Adhya TK (2009) Methane and nitrous oxide emissions from an integrated rainfed rice–fish farming system of Eastern India. *Agriculture, Ecosystems & Environment*, **129**, 228–237.
- DeFries R, Rosenzweig C (2010) Toward a whole-landscape approach for sustainable land use in the tropics. *Proceedings of the National Academy of Sciences of the United States of America*, **107**, 19627–19632.
- Deininger K, Byerlee D (2011). *Rising Global Interest in Farmland: Can it Yield Sustainable and Equitable Benefits?* Agriculture and Rural Development series, The World Bank, Washington, DC.
- Ellert BH, Janzen HH (2008) Nitrous oxide, carbon dioxide and methane emissions from irrigated cropping systems as influenced by legumes, manure and fertilizer. *Canadian Journal of Soil Science*, **88**, 207–217.
- Firestone MK, Davidson EA (1989) Microbial basis of NO and N₂O production and consumption in soils. In: *Exchange of Trace Gases Between Terrestrial Ecosystems and the Atmosphere* (eds Andreae MO, Schimel DS), pp. 7–21. John Wiley & Sons, New York.
- Garnett T (2009) Livestock-related greenhouse gas emissions: impacts and options for policy makers. *Environmental Science & Policy*, **12**, 491–503.
- Ghosh S, Majumdar D, Jain MC (2003) Methane and nitrous oxide emissions from an irrigated rice of North India. *Chemosphere*, **51**, 181–195.
- Godfray HCJ, Pretty J, Thomas SM, Warham EJ, Beddington JR (2011) Linking policy on climate and food. *Science*, **331**, 1013–1014.
- Grandy AS, Loeckle TD, Parr S, Robertson GP (2006) Long-term trends in nitrous oxide emissions, soil nitrogen, and crop yields of till and no-till cropping systems. *Journal of Environmental Quality*, **35**, 1487–1495.
- van Groenigen JW, Velthof GL, Oenema O, van Groenigen KJ, van Kessel C (2010) Towards an agronomic assessment of N₂O emissions: a case study for arable crops. *European Journal of Soil Science*, **61**, 903–913.
- Guo ZL, Cai CF, Li ZX, Wang TW, Zheng MJ (2009) Crop residue effect on crop performance, soil N₂O and CO₂ emissions in alley cropping systems in subtropical China. *Agroforestry Systems*, **76**, 67–80.
- Halvorson AD, Del Grosso SJ, Alluvione F (2010) Nitrogen source effects on nitrous oxide emissions from irrigated no-till corn. *Journal of Environmental Quality*, **39**, 1554–1562.
- Hedges LV, Gurevitch J, Curtis PS (1999) The meta-analysis of response ratios in experimental ecology. *Ecology*, **80**, 1150–1156.
- Hill J, Nelson E, Tilman D, Polasky S, Tiffany D (2006) Environmental, economic, and energetic costs and benefits of biodiesel and ethanol biofuels. *Proceedings of the National Academy of Sciences of the United States of America*, **103**, 11206–11210.
- Hoben JP, Gehl RJ, Millar N, Grace PR, Robertson GP (2011) Nonlinear nitrous oxide (N₂O) response to nitrogen fertilizer in on-farm corn crops of the US Midwest. *Global Change Biology*, **17**, 1140–1152.
- Holzappel-Pschorn A, Seiler W (1986) Methane emission during a cultivation period from an Italian rice paddy. *Journal of Geophysical Research*, **91**, 11803–11814.
- Holzappel-Pschorn A, Conrad R, Seiler W (1986) Effects of vegetation on the emission of methane from submerged paddy soil. *Plant and Soil*, **92**, 223–233.
- Hou AX, Chen GX, Wang ZP, Van Cleemput O, Patrick WH (2000) Methane and nitrous oxide emissions from a rice field in relation to soil redox and microbiological processes. *Soil Science Society of America Journal*, **64**, 2180–2186.
- Hütsch BW (2001) Methane oxidation in non-flooded soils as affected by crop production – invited paper. *European Journal of Agronomy*, **14**, 237–260.
- IPCC (1995) Other trace gases and atmospheric chemistry. In: *Climate Change 1994: Radiative Forcing of Climate Change and an Evaluation of the IPCC IS92 Emission Scenarios*, (eds Houghton JT, Meira Filho LG, Bruce J, Lee H, Callander BA, Harris N, Maskell K), pp. 73–127. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- IPCC (2001) Radiative forcing of climate change. In: *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, (eds Houghton JT, Ding Y, Griggs DJ, Noguer M, van der Linden PJ, Dai X, Maskell K, Johnson CA), 881 pp. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- IPCC (2006) Agriculture, forestry and other land use. In: *2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme*, (eds Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K), 66 pp. Institute for Global Environmental Strategies, Hayama, Japan.
- Jayasundara S, Wagner-Riddle C, Parkin G, Von Bertoldi P, Warland J, Kay B, Voroney P (2007) Minimizing nitrogen losses from a corn–soybean–winter wheat rotation with best management practices. *Nutrient Cycling in Agroecosystems*, **79**, 141–159.
- Ju XT, Xing GX, Chen XP *et al.* (2009) Reducing environmental risk by improving N management in intensive Chinese agricultural systems. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 3041–3046.
- Kaiser EA, Kohrs K, Kucke M, Schnug E, Heinemeyer O, Munch JC (1998) Nitrous oxide release from arable soil: importance of N-fertilization, crops and temporal variation. *Soil Biology & Biochemistry*, **30**, 1553–1563.
- Kessavalou A, Mosier AR, Doran JW, Drijber RA, Lyon DJ, Heinemeyer O (1998) Fluxes of carbon dioxide, nitrous oxide, and methane in grass sod and winter wheat–fallow tillage management. *Journal of Environmental Quality*, **27**, 1094–1104.
- Khalil MI, Rosenani AB, Van Cleemput O, Fauziah CI, Shamsuddin J (2002) Nitrous oxide emissions from an ultisol of the humid tropics under maize–groundnut rotation. *Journal of Environmental Quality*, **31**, 1071–1078.
- Kreye C, Dittert K, Zheng X, Zhang X, Lin S, Tao H, Sattelmacher B (2007) Fluxes of methane and nitrous oxide in water-saving rice production in north China. *Nutrient Cycling in Agroecosystems*, **77**, 293–304.
- Li X, Zhang X, Xu H, Cai Z, Yagi K (2009) Methane and nitrous oxide emissions from rice paddy soil as influenced by timing of application of hydroquinone and dicyandiamide. *Nutrient Cycling in Agroecosystems*, **85**, 31–40.
- Lindau CW, Bollich PK, Delaune RD, Patrick WH, Law VJ (1991) Effect of urea fertilizer and environmental factors on CH₄ emissions from a Louisiana, USA rice field. *Plant and Soil*, **136**, 195–203.
- Liu C, Wang K, Meng S *et al.* (2011) Effects of irrigation, fertilization and crop straw management on nitrous oxide and nitric oxide emissions from a wheat–maize rotation field in northern China. *Agriculture, Ecosystems & Environment*, **140**, 226–233.
- Lobell DB, Cassman KG, Field CB (2009) Crop yield gaps: their importance, magnitudes, and causes. *Annual Review of Environment and Resources*, **34**, 179–204.

- Ma J, Li XL, Xu H, Han Y, Cai ZC, Yagi K (2007) Effects of nitrogen fertilizer and wheat straw application on CH₄ and N₂O emissions from a paddy rice field. *Australian Journal of Soil Research*, **45**, 359–367.
- Ma J, Ma E, Xu H, Yagi K, Cai Z (2009) Wheat straw management affects CH₄ and N₂O emissions from rice fields. *Soil Biology & Biochemistry*, **41**, 1022–1028.
- Ma BL, Wu TY, Tremblay N *et al.* (2010a) Nitrous oxide fluxes from corn fields: on-farm assessment of the amount and timing of nitrogen fertilizer. *Global Change Biology*, **16**, 156–170.
- Ma E, Zhang G, Ma J, Xu H, Cai Z, Yagi K (2010b) Effects of rice straw returning methods on N₂O emission during wheat-growing season. *Nutrient Cycling in Agroecosystems*, **88**, 463–469.
- Maclean JL, Dawe DC, Hardy B, Hettel GP (eds) (2002) *Rice Almanac. Source Book for the Most Important Economic Activity on Earth* (3rd edn). CABI Publishing, Wallingford, UK.
- Majumdar D, Pathak H, Kumar S, Jain MC (2002) Nitrous oxide emission from a sandy loam inceptisol under irrigated wheat in India as influenced by different nitrification inhibitors. *Agriculture, Ecosystems & Environment*, **91**, 283–293.
- Malhi SS, Lemke R (2007) Tillage, crop residue and N fertilizer effects on crop yield, nutrient uptake, soil quality and nitrous oxide gas emissions in a second 4-yr rotation cycle. *Soil & Tillage Research*, **96**, 269–283.
- Malla G, Bhatia A, Pathak H, Prasad S, Jain N, Singh J (2005) Mitigating nitrous oxide and methane emissions from soil in rice-wheat system of the Indo-Gangetic plain with nitrification and urease inhibitors. *Chemosphere*, **58**, 141–147.
- Masscheleyn PH, Delaune RD, Patrick WH Jr (1993) Methane and nitrous oxide emissions from laboratory measurements of rice soil suspension: effect of soil oxidation-reduction status. *Chemosphere*, **26**, 251–260.
- Matson PA, Parton WJ, Power AG, Swift MJ (1997) Agricultural intensification and ecosystem properties. *Science*, **277**, 504–509.
- McSwiney CP, Robertson GP (2005) Nonlinear response of N₂O flux to incremental fertilizer addition in a continuous maize (*Zea mays* L.) cropping system. *Global Change Biology*, **11**, 1712–1719.
- Mosier A, Wassmann R, Verchot L, King J, Palm C (2004) Methane and nitrogen oxide fluxes in tropical agricultural soils: sources, sinks and mechanisms. *Environment, Development and Sustainability*, **6**, 11–49.
- Mosier AR, Halvorson AD, Reule CA, Liu XJ (2006) Net global warming potential and greenhouse gas intensity in irrigated cropping systems in Northeastern Colorado. *Journal of Environmental Quality*, **35**, 1584–1598.
- Pan G, Xu X, Smith P, Pan W, Lal R (2010) An increase in topsoil SOC stock of China's croplands between 1985 and 2006 revealed by soil monitoring. *Agriculture, Ecosystems & Environment*, **136**, 133–138.
- Parkin TB, Hatfield JL (2010) Influence of nitrapyrin on N₂O losses from soil receiving fall-applied anhydrous ammonia. *Agriculture, Ecosystems & Environment*, **136**, 81–86.
- Parkin TB, Kaspar TC (2006) Nitrous oxide emissions from corn–soybean systems in the Midwest. *Journal of Environmental Quality*, **35**, 1496–1506.
- Pathak H, Bhatia A, Prasad S, Singh S, Kumar S, Jain MC, Kumar U (2002) Emission of nitrous oxide from rice-wheat systems of Indo-Gangetic plains of India. *Environmental Monitoring and Assessment*, **77**, 163–178.
- Pathak H, Singh R, Bhatia A, Jain N (2006) Recycling of rice straw to improve wheat yield and soil fertility and reduce atmospheric pollution. *Paddy and Water Environment*, **4**, 111–117.
- Pathak H, Jain N, Bhatia A, Patel J, Aggarwal PK (2010) Carbon footprints of Indian food items. *Agriculture, Ecosystems & Environment*, **139**, 66–73.
- Phillips RL, Tanaka DL, Archer DW, Hanson JD (2009) Fertilizer application timing influences greenhouse gas fluxes over a growing season. *Journal of Environmental Quality*, **38**, 1569–1579.
- Post WM, Izaurralde RC, Mann LK, Bliss N (2001) Monitoring and verifying changes of organic carbon in soil. *Climatic Change*, **51**, 73–99.
- Qian JH, Doran JW, Weier KL, Mosier AR, Peterson TA, Power JF (1997) Soil denitrification and nitrous oxide losses under corn irrigated with high-nitrate groundwater. *Journal of Environmental Quality*, **26**, 348–360.
- Qin Y, Liu S, Guo Y, Liu Q, Zou J (2010) Methane and nitrous oxide emissions from organic and conventional rice cropping systems in Southeast China. *Biology and Fertility of Soils*, **46**, 825–834.
- Robertson GP, Grace PR (2004) Greenhouse gas fluxes in tropical and temperate agriculture: the need for a full-cost accounting of global warming potentials. *Environment, Development and Sustainability*, **6**, 51–63.
- Robertson GP, Vitousek PM (2009) Nitrogen in agriculture: balancing the cost of an essential resource. *Annual Review of Environment and Resources*, **34**, 97–125.
- Robertson GP, Paul EA, Harwood RR (2000) Greenhouse gases in intensive agriculture: contributions of individual gases to the radiative forcing of the atmosphere. *Science*, **289**, 1922–1925.
- Rosenberg MS, Adams DC, Gurevitch J (2000) *METAWIN, Statistical Software for Meta-Analysis*, Version 2. Sinauer, Sunderland, MA.
- Ruser R, Flessa H, Schilling R, Beese F, Munch JC (2001) Effect of crop-specific field management and N fertilization on N₂O emissions from a fine-loamy soil. *Nutrient Cycling in Agroecosystems*, **59**, 177–191.
- Sachs J, Remans R, Smukler S *et al.* (2010) Monitoring the world's agriculture. *Nature*, **466**, 558–560.
- Sass RL, Fisher FM, Harcombe PA, Turner FT (1990) Methane production and emission in a Texas rice field. *Global Biogeochemical Cycles*, **4**, 47–68.
- Sehy U, Ruser R, Munch JC (2003) Nitrous oxide fluxes from maize fields: relationship to yield, site-specific fertilization, and soil conditions. *Agriculture, Ecosystems & Environment*, **99**, 97–111.
- Shang Q, Yang X, Gao C *et al.* (2011) Net annual global warming potential and greenhouse gas intensity in Chinese double rice-cropping systems: a 3-year field measurement in long-term fertilizer experiments. *Global Change Biology*, **17**, 2196–2210.
- Six J, Ogle SM, Breidt FJ, Conant RT, Mosier AR, Paustian K (2004) The potential to mitigate global warming with no-tillage management is only realized when practised in the long term. *Global Change Biology*, **10**, 155–160.
- Smith P, Martino D, Cai Z *et al.* (2007) *Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Metz B, Davidson OR, Bosch PR, Dave R, Meyer LA), pp. 497–540. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- Steffan-Dewenter I, Kessler M, Barkmann J *et al.* (2007) Tradeoffs between income, biodiversity, and ecosystem functioning during tropical rainforest conversion and agroforestry intensification. *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 4973–4978.
- Sterling J, Buttner P (2009) *Wildlife known to use California rielands*. California Rice Commission. ICF Jones & Stokes, Sacramento, CA.
- Stewart CE, Paustian K, Conant RT, Plante AF, Six J (2007) Soil carbon saturation: concept, evidence and evaluation. *Biogeochemistry*, **86**, 19–31.
- Thenkabail PS, Hanjra MA, Dheeravath V, Gumma M (2010) A holistic view of global croplands and their water use for ensuring global food security in the 21st century through advanced remote sensing and non-remote sensing approaches. *Remote Sensing*, **2**, 211–261.
- Tilman D (1999) Global environmental impacts of agricultural expansion: the need for sustainable and efficient practices. *Proceedings of the National Academy of Sciences of the United States of America*, **96**, 5995–6000.
- Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S (2002) Agricultural sustainability and intensive production practices. *Nature*, **418**, 671–677.
- Towprayoon S, Smakgahn K, Poonkaew S (2005) Mitigation of methane and nitrous oxide emissions from drained irrigated rice fields. *Chemosphere*, **59**, 1547–1556.
- Venterea RT, Dolan MS, Ochsner TE (2010) Urea decreases nitrous oxide emissions compared with anhydrous ammonia in a Minnesota corn cropping system. *Soil Science Society of America Journal*, **74**, 407–418.
- Vitousek PM, Mooney HA, Lubchenco J, Melillo JM (1997) Human domination of earth's ecosystems. *Science*, **277**, 494–499.
- Wagner-Riddle C, Furon A, Mclaughlin NL *et al.* (2007) Intensive measurement of nitrous oxide emissions from a corn–soybean–wheat rotation under two contrasting management systems over 5 years. *Global Change Biology*, **13**, 1722–1736.
- Wassmann R, Lantin RS, Neue HU, Buendia LV, Corton TM, Lu Y (2000) Characterization of methane emissions from rice fields in Asia. III. Mitigation options and future research needs. *Nutrient Cycling in Agroecosystems*, **58**, 23–36.
- Webb J, Ellis S, Harrison R, Thorman R (2004) Measurement of N fluxes and soil N in two arable soils in the UK. *Plant and Soil*, **260**, 253–270.
- Wei XR, Hao MD, Xue XH, Shi P, Horton R, Wang A, Zang YF (2010) Nitrous oxide emission from highland winter wheat field after long-term fertilization. *Biogeochemistry*, **7**, 3301–3310.
- Weiske A, Benckiser G, Herbert T, Ottow JCG (2001) Influence of the nitrification inhibitor 3,4-dimethylpyrazole phosphate (DMPP) in comparison to dicyandiamide (DCD) on nitrous oxide emissions, carbon dioxide fluxes and methane oxidation during 3 years of repeated application in field experiments. *Biology and Fertility of Soils*, **34**, 109–117.
- West TO, Post WM (2002) Soil organic carbon sequestration rates by tillage and crop rotation: a global data analysis. *Soil Science Society of America Journal*, **66**, 1930–1946.
- Wu J (2011) Carbon accumulation in paddy ecosystems in subtropical China: evidence from landscape studies. *European Journal of Soil Science*, **62**, 29–34.
- Xie B, Zheng X, Zhou Z *et al.* (2010) Effects of nitrogen fertilizer on CH₄ emission from rice fields: multi-site field observations. *Plant and Soil*, **326**, 393–401.

- Yagi K, Tsuruta H, Minami K (1997) Possible options for mitigating methane emission from rice cultivation. *Nutrient Cycling in Agroecosystems*, **49**, 213–220.
- Yan X, Hosen Y, Yagi K (2001) Nitrous oxide and nitric oxide emissions from maize field plots as affected by N fertilizer type and application method. *Biology and Fertility of Soils*, **34**, 297–303.
- Yan X, Akiyama H, Yagi K, Akimoto H (2009) Global estimations of the inventory and mitigation potential of methane emissions from rice cultivation conducted using the 2006 Intergovernmental Panel on Climate Change Guidelines. *Global Biogeochemical Cycles*, **23**, GB2002.
- Yao Z, Zhou Z, Zheng X *et al.* (2010) Effects of organic matter incorporation on nitrous oxide emissions from rice-wheat rotation ecosystems in China. *Plant and Soil*, **327**, 315–330.
- Yu KW, Wang ZP, Chen GX (1997) Nitrous oxide and methane transport through rice plants. *Biology and Fertility of Soils*, **24**, 341–343.
- Zhang A, Cui L, Pan G *et al.* (2010) Effect of biochar amendment on yield and methane and nitrous oxide emissions from a rice paddy from Tai Lake plain, China. *Agriculture, Ecosystems & Environment*, **139**, 469–475.
- Zou J, Huang Y, Jiang J, Zheng X, Sass RL (2005) A 3-year field measurement of methane and nitrous oxide emissions from rice paddies in China: Effects of water regime, crop residue, and fertilizer application. *Global Biogeochemical Cycles*, **19**, GB2021.
- Zou J, Huang Y, Zheng X, Wang Y (2007) Quantifying direct N₂O emissions in paddy fields during rice growing season in mainland China: dependence on water regime. *Atmospheric Environment*, **41**, 8030–8042.