

# Seasonal Methane and Nitrous Oxide Emissions of Several Rice Cultivars in Direct-Seeded Systems

Maegen B. Simmonds,\* Merle Anders, Maria Arlene Adviento-Borbe, Chris van Kessel, Anna McClung, and Bruce A. Linquist

## Abstract

An understanding of cultivar effects on field greenhouse gas (GHG) emissions in rice (*Oryza sativa* L.) systems is needed to improve the accuracy of predictive models used for estimating GHG emissions and to evaluate the GHG mitigation potential of different cultivars. We compared CH<sub>4</sub> and N<sub>2</sub>O emissions, global warming potential (GWP = N<sub>2</sub>O + CH<sub>4</sub>), yield-scaled GWP (GWP<sub>y</sub> = GWP Mg<sup>-1</sup> grain), and plant growth characteristics of eight cultivars within four study sites in California and Arkansas. Nitrous oxide emissions were negligible (<10% of GWP) and were not different among cultivars. Seasonal CH<sub>4</sub> emissions differed between cultivars by a factor of 2.1 and 1.4 at one California and one Arkansas site, respectively. Plant growth characteristics were generally not correlated with seasonal CH<sub>4</sub> emissions; however, the strongest correlations were observed for shoot and total plant (root + shoot) biomass at heading ( $r = 0.60$ ) at one California site and for grain at maturity ( $r = -0.95$ ) at one Arkansas site. Although differences in GWP and GWP<sub>y</sub> were observed, there were inconsistencies across sites, indicating the importance of the genotype × environment interaction. Overall, the cultivars with the lowest CH<sub>4</sub> emissions, GWP, and GWP<sub>y</sub> at the California and Arkansas sites were the lowest and highest yielding, respectively. These findings highlight the potential for breeding high-yielding cultivars with low GWP, the ideal scenario to achieve low GWP<sub>y</sub>, but environmental conditions must also be considered.

AGRICULTURE is a major source of global anthropogenic emissions of greenhouse gases (GHGs), accounting for an estimated 47 and 58% of CH<sub>4</sub> and N<sub>2</sub>O emissions, respectively (Smith et al., 2007). With a radiative forcing capacity of 25 and 298 times greater than CO<sub>2</sub> on a 100-yr timescale, respectively, their unprecedented increase in atmospheric concentrations has led to international efforts to measure and mitigate emissions (Forster et al., 2007). The cultivation of rice (*Oryza sativa* L.) is an important source of global anthropogenic CH<sub>4</sub> emissions, with estimates ranging from 25.6 Tg CH<sub>4</sub> yr<sup>-1</sup> (Yan et al., 2009) to 112 Tg CH<sub>4</sub> yr<sup>-1</sup> (Chen and Prinn, 2006). Although agricultural soils in general contribute an estimated 5.6 Tg N<sub>2</sub>O yr<sup>-1</sup> (Smith et al., 2007), the vast majority (89%) of the global warming potential (GWP) of rice systems is due to CH<sub>4</sub> emissions (Linquist et al., 2012). Compared with other major cereal crops (e.g., wheat [*Triticum aestivum* L.] and maize [*Zea mays* L.]), the GWP of CH<sub>4</sub> and N<sub>2</sub>O emissions from rice production is on average approximately four times greater (Linquist et al., 2012).

When rice fields are flooded, N<sub>2</sub>O fluxes from soil remain low, and the primary pathway of N<sub>2</sub>O emissions is via transport through the intracellular air spaces within the rice plants (aerenchyma cells), which comprises, on average, 80% (Yu et al., 1997) to 87% (Yan et al., 2000) of total N<sub>2</sub>O emissions. During nonflooded conditions, a considerably lower proportion (18%) of total N<sub>2</sub>O is emitted through the rice plant (Yan et al., 2000). Rice plants also regulate CH<sub>4</sub> emissions by facilitating its production, oxidation, and transport (Wassmann and Aulakh, 2000). Microbial decomposition of organic matter in flooded, oxygen-depleted soils is the process by which methanogenesis occurs (Conrad, 2005). Rice plants influence CH<sub>4</sub> production by supplying readily decomposable root-derived C substrates to methanogenic bacteria in the form of exudates, sloughed cells, mucilage, and litter (Lu et al., 2000b, 2002; Aulakh et al., 2001b). Methane transport from the soil to the atmosphere via aerenchyma cells of the rice plant accounts for the vast majority

Copyright © American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America. 5585 Guilford Rd., Madison, WI 53711 USA. All rights reserved. No part of this periodical may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher.

J. Environ. Qual. 44:103–114 (2015)

doi:10.2134/jeq2014.07.0286

Received 3 July 2014.

Accepted 30 Oct. 2014.

\*Corresponding author (mbsimmonds@ucdavis.edu).

M.B. Simmonds, M.A. Adviento-Borbe, C. van Kessel, and B.A. Linquist Dep. of Plant Sciences, Univ. of California, Davis, One Shields Avenue, Davis, CA 95616; M. Anders, Univ. of Arkansas Rice Research & Extension Center, Stuttgart, AR 72160; A. McClung, Dale Bumpers National Rice Research Center, Stuttgart, AR 72160. Assigned to Associate Editor Pierre-Andre Jacinthe.

**Abbreviations:** GC, gas chromatograph; GHG, greenhouse gas; GWP, global warming potential; GWP<sub>y</sub>, yield-scaled global warming potential; SOC, soil organic carbon.

of CH<sub>4</sub> emissions (range of estimates, 80–98%); molecular diffusion through surface irrigation water and ebullition are secondary sources of CH<sub>4</sub> emissions (Holzapfel-Pschorn et al., 1986; Holzapfel-Pschorn and Seiler, 1986; Tyler et al., 1997; Yu et al., 1997). The aerenchyma cells also serve as a conduit for transport of O<sub>2</sub> from the atmosphere to the roots, providing a sufficient aerobic environment for root respiration in flooded soil conditions (Colmer, 2003; Suralta and Yamauchi, 2008) and, in turn, supporting CH<sub>4</sub> oxidation in the rhizosphere (Sigren et al., 1997; Eller and Frenzel, 2001). Some studies have shown differences in CH<sub>4</sub> emissions among cultivars due to varying degrees of controls over these mechanisms (Huang et al., 1997; Bilek et al., 1999). In a 2-yr study, Sigren et al. (1997) showed that two Texas cultivars differed in CH<sub>4</sub> emissions and attributed these differences in C substrate availability in the root zone and to the lack of differences in gas transport capacity. In contrast, Butterbach-Bahl et al. (1997) deduced that differences in CH<sub>4</sub> transport capacity were driving the observed differences in CH<sub>4</sub> emissions between two Italian cultivars because neither CH<sub>4</sub> oxidation nor CH<sub>4</sub> production differed between cultivars.

There is also evidence that cultivar differences in CH<sub>4</sub> emissions are not persistent over time. In a 7-yr field study, Wassmann et al. (2002) reported cultivar differences in CH<sub>4</sub> emissions in some seasons, but, over the course of nine growing seasons, no consistent distinctions among cultivars were found. Rather, the cultivars displayed high phenotypic plasticity due to complex interactions with their environment. Similarly, Lu et al. (2000a) found that the differences between hybrid and conventional cultivars in CH<sub>4</sub> emissions depended on the season.

Specific mechanisms for cultivar differences in CH<sub>4</sub> emissions have been related to plant growth and morphological characteristics. Aulakh et al. (2000) found that CH<sub>4</sub> transport capacity of 12 rice cultivars generally increased during the vegetative growth period, after which three divergent patterns in CH<sub>4</sub> transport capacity occurred; in hybrids it gradually increased to maturity, and in inbred cultivars it either rapidly declined from flowering to maturity or changed minimally. These distinctions in patterns of CH<sub>4</sub> transport capacity corresponded to root and shoot biomass during the vegetative growth period and to total root, shoot, and grain biomass from panicle initiation to maturity. These relationships were attributed to differences in the duration of radial development of aerenchyma and total number of outlets (tillers) of aerenchyma. Positive correlations have also been demonstrated between rates of root exudation and root, shoot, and grain biomass (Aulakh et al., 2001a) and between CH<sub>4</sub> production from root exudates and root and shoot biomass (Aulakh et al., 2001b).

With the significant contribution of rice cultivation to global CH<sub>4</sub> emissions and projections for increased rice production to meet global food demand, a clear challenge exists to improve yields while minimizing CH<sub>4</sub> emissions from rice production (Smith et al., 2007). Whereas soil, climate, and management factors are accounted for in many process-based models, cultivar-specific mechanisms are not. Although many of these models have the capacity to specify basic crop growth parameters, there is limited cultivar-specific data available for improving the scientific structure and calibration of these models, particularly in the rice growing regions in California and the southern U.S. states. A new understanding of the magnitude of cultivar differences is critical

for accurate modeling of GHG emissions from rice systems, which is of growing importance due to emerging markets for C trading in the agricultural sector (Climate Action Reserve, 2013).

We conducted multi-year field studies in California and Arkansas. Our main objectives were (i) to determine if seasonal CH<sub>4</sub> and N<sub>2</sub>O emissions, GWP, and yield-scaled GWP (GWP<sub>Y</sub> = GWP Mg<sup>-1</sup> grain) differ among several commonly grown cultivars in California and the southern United States and (ii) to determine the relative importance of plant growth characteristics in explaining the variability in seasonal GHG emissions. Our overall hypothesis was that cultivars differ in CH<sub>4</sub> emissions and have distinct plant growth characteristics related to differences in magnitude of emissions but do not differ in N<sub>2</sub>O emissions due to minimal seasonal N<sub>2</sub>O fluxes during the continuously flooded field conditions. Thus, cultivar GWP was predicted to differ according to differences primarily in CH<sub>4</sub> emissions. Specifically, we expected cultivars with the highest root and shoot biomass to have the highest GWP due to a higher capacity for CH<sub>4</sub> production via greater root exudation and to have a higher capacity for CH<sub>4</sub> transport via a larger, or a more extensive, aerenchyma system. Among cultivars with similar root and shoot biomass, those with higher grain yield were expected to have lower CH<sub>4</sub> emissions due to greater allocation of C to grain versus root exudation. Therefore, we predicted high-yielding cultivars with low root and shoot biomass at heading to have the lowest GWP<sub>Y</sub>.

## Materials and Methods

### Field Experiment

Experiments were conducted in experimental fields in California and Arkansas during the 2011 and 2012 rice-growing seasons. In California, field sites were located at the Rice Experiment Station (CA-1) in Biggs in 2011 (39°27'47'' N, 121°43'35'' W) and at the University of California, Davis (CA-2) in 2012 (38°32'33.20'' N, 121°48'38.69'' W). In Arkansas, field sites were located at the University of Arkansas Rice Research and Extension Center near Stuttgart, AR, in 2011 (AR-1) and 2012 (AR-2) (34°28'8.64'' N, 91°25'7.19'' W and 34°27'54.6942'' N, 91°24'21.8484'' W, respectively). Soils at CA-1 and CA-2 are classified as Esquon-Neerdobe complex (Fine, smectitic, thermic Xeric Epiaquerts and Duraquerts) and Capay silty clay (Fine, smectitic, thermic Typic Haploxerert), respectively, and soils at AR-1 and AR-2 are classified as Dewitt silt loam (Fine, smectitic, thermic, Typic Albaqualfs).

Eight cultivars adapted to the California and southern U.S. environments were investigated for differences in seasonal CH<sub>4</sub> and N<sub>2</sub>O emissions (Table 1). All of the cultivars used are commercially produced for conventional or specialty markets and thus are a representation of material from current varietal improvement programs. The field trials were arranged in a randomized complete block design, with three replications in CA-1, CA-2, and AR-2 and only one replication in AR-1.

Standard management practices for water-seeded and drill-seeded systems were followed at the California and Arkansas sites, respectively (Table 2). Plot sizes were 4.6 and 20 m<sup>2</sup> at CA-1 and CA-2, respectively, and 24 m<sup>2</sup> at AR-1 and AR-2.

**Table 1. Description of cultivars tested and the locations of the field experiments.**

Cultivar	Environment adapted to	Description	Market class	Study site†
Calmati-202	California	specialty	long-grain	CA-1, CA-2
L206	California	conventional	long-grain	CA-1, CA-2
M206	California	conventional	medium-grain	CA-1, CA-2
CLXL745	southern U.S.	hybrid	long-grain	CA-2, AR-2
CLXP4534	southern U.S.	hybrid	long-grain	CA-2, AR-1
Francis	southern U.S.	conventional	long-grain	AR-1, AR-2
Jupiter	southern U.S.	conventional	medium-grain	AR-1, AR-2
Sabine	southern U.S.	specialty	long-grain	AR-1, AR-2

† AR-1 and AR-2 refer to Arkansas field sites located at the University of Arkansas Rice Research and Extension Center near Stuttgart, AR, in 2011 and 2012, respectively; CA-1 and CA-2 refer to California field sites located at the Rice Experiment Station in Biggs in 2011 and at the University of California, Davis in 2012, respectively.

**Table 2. Summary of management practices at the four study sites.**

Management practice	Study site†			
	CA-1	CA-2	AR-1	AR-2
Previous year's crop	fallow	fallow for 4 yr	soybean	soybean
Residue management	crop residue burned	only weeds incorporated in soil	crop residue incorporated in soil	crop residue incorporated in soil
Water management	no winter flood	no winter flood	no winter flood	no winter flood
Method of direct seeding	water-seeded	water-seeded	drill-seeded	drill-seeded
Seeding rate, kg ha <sup>-1</sup>	168	123 for non-hybrids, 50 for hybrids	123 for non-hybrids, 28 for hybrids	123 for non-hybrids, 28 for hybrids
Date of seeding	27 May 2011	4 June 2012	17 May 2011	11 Apr. 2012
Water management	continuous flood	continuous flood	one flush irrigation before continuous flood	continuous flood
Date of permanent flood	25 May 2011	18 May 2012	20 June 2011	17 May 2012
Date of drain	23 Sept. 2011	8 Oct. 2012	11 Sept. 2011	18 Aug. 2012
Fertility management				
Date of fertilization and fertilizer type	21 May 2011 (urea blend, 34:17:0) and 11 July 2011 [(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> ]	17 May 2012 [Ca(H <sub>2</sub> PO <sub>4</sub> ) <sub>2</sub> ·H <sub>2</sub> O and K <sub>2</sub> SO <sub>4</sub> ], 18 May 2012 (urea), 6 July 2012 (urea)	early spring pre-tillage 2011 [Ca(H <sub>2</sub> PO <sub>4</sub> ) <sub>2</sub> ·H <sub>2</sub> O and KCl], 20 June 2011 (urea)	early spring pre-tillage 2012 [Ca(H <sub>2</sub> PO <sub>4</sub> ) <sub>2</sub> ·H <sub>2</sub> O and KCl], 16 May 2012 (urea)
Total N, kg ha <sup>-1</sup>	130	210	168	134
First application	95	160	168	134
Second application	30	50		
Total P, kg ha <sup>-1</sup>	21	24	29	29
Total K, kg ha <sup>-1</sup>	0	46	84	84
Date of harvest	24 Oct. 2011	29 Oct. 2012	26 Sept. 2011	28 Aug. 2012

† AR-1 and AR-2 refer to Arkansas field sites located at the University of Arkansas Rice Research and Extension Center near Stuttgart, AR, in 2011 and 2012, respectively; CA-1 and CA-2 refer to California field sites located at the Rice Experiment Station in Biggs in 2011 and at the University of California, Davis in 2012, respectively.

All sites were planted within the optimal planting dates for the region. Seeding rates and fertility management were based on a combination of considerations: recommendations for the region and type of seeding system, knowledge of the field site, and the specific cultivars tested. Due to failure of the first seeding on 18 May 2012 to germinate at CA-2, the plots were reseeded in the same field on 4 June 2012.

## Soil Sampling

After spring tillage operations were completed and before seeding and flooding, soil samples were collected from 0 to 15 cm depth from each field to characterize soil chemical and physical properties at each site. Soil was analyzed for soil texture and various chemical properties (pH, electrical conductivity, total organic C, total N, extractable P, and exchangeable K) using standard methods (Table 3).

## Gas Sampling and Analysis

Daily fluxes of CH<sub>4</sub> and N<sub>2</sub>O were measured using a vented closed-chamber method (Hutchinson and Livingston, 1993) with four sampling times (0, 21, 42, 63 min.). The construction and installation of the chambers followed the methods described by Adviento-Borbe et al. (2013). Gas sampling began within 5 d of the onset of the permanent flood at the CA sites and at 22 and 12 d after the onset of the permanent flood at AR-1 and AR-2, respectively, and continued until harvest or within 4 d before harvest. The 63-min gas sampling period occurred between 0900 and 1400 h on a weekly basis at CA-2, AR-1, and AR-2; biweekly at CA-1; and up to daily during drainage or flooding events at all sites. Two chambers were installed at CA-2, AR-1, and AR-2 in each plot to reduce disturbance to the rice plants during sampling by alternating which chambers were used. At CA-1, one chamber was installed per plot due to less frequent sampling.

To determine concentrations of CH<sub>4</sub> and N<sub>2</sub>O in the chamber headspace, gas samples were analyzed using a GC-2014 gas chromatograph (GC) (Shimadzu Scientific, Inst.) with a flame ionization detector for CH<sub>4</sub> and a <sup>63</sup>Ni electron capture detector for N<sub>2</sub>O. Detection limits of the GC instrument were 2.2 pg s<sup>-1</sup> for CH<sub>4</sub> and 0.3 pg s<sup>-1</sup> for N<sub>2</sub>O based on the results given by the Shimadzu technical engineers who conducted the QC/QA of the instrument and corresponding packed columns for ambient GHG detections. We also calculated the flux chamber detection limits based on (i) four gas levels with time zero, (ii) chamber closure of 63 min, and (iii) linear flux equation (Parkin et al., 2012); they were 0.00076 ppm<sub>v</sub> N<sub>2</sub>O min<sup>-1</sup> and 0.0072 ppm<sub>v</sub> CH<sub>4</sub> min<sup>-1</sup> for three time intervals. However, the detection limit of the GC was higher (less sensitive), and therefore those detection limits were used. Using the ideal gas law and measurements of air temperature and volume within the chambers, headspace gas concentrations were converted to mass per volume (g N<sub>2</sub>O or CH<sub>4</sub> L<sup>-1</sup>). Fluxes of CH<sub>4</sub> and N<sub>2</sub>O were estimated from the slope of the linear regression of gas concentration by chamber closure time divided by the surface area within the chamber. Diurnal variation in GHG emissions was not accounted for in daily flux calculations based on the assumption that soil temperature represented average daily values during the period that gas sampling occurred (i.e., 0900–1400 h) (Bossio et al., 1999). Fluxes were set to zero if the change in emissions was less than the minimum detection limit of the GC. For quality assurance of the non-zero flux estimates, we used the distribution of all the RMSE values of the flux regressions as criteria for assessing the reliability of the data (i.e., whether there was excessive scatter about a particular regression, potentially due to measurement error, or a reasonable amount of scatter relative to the entirety of the dataset). Accordingly, if the RMSE of the linear regression of gas concentration by chamber closure time was >2 SD above the mean RMSE of all the linear regressions at a given site, the corresponding flux value was treated as an outlier, and the value was omitted from further analysis.

Cumulative seasonal emissions were estimated using linear interpolation of the daily fluxes. Total GWP of seasonal CH<sub>4</sub> and N<sub>2</sub>O emissions was calculated in units of CO<sub>2</sub> equivalents (CO<sub>2</sub> eq) with a 100-yr time frame using radiative forcing potentials of 25 and 298 relative to CO<sub>2</sub>, respectively (Forster et al., 2007). Yield-scaled GWP is the ratio of GWP to the mass of grain yield (i.e., kg CO<sub>2</sub> eq Mg<sup>-1</sup>). One GWP data point and the corresponding GWP<sub>Y</sub> data point (for cultivar CLXL745, rep 3) were negative and therefore were treated as outliers and omitted.

## Plant Sampling

Uniform stand areas within each plot were used for plant sampling. Root and shoot biomass were measured at the heading stage (exertion of the panicle) of the rice plants at all sites except AR-1. Root and shoot biomass per unit area was estimated from a 0.1-m<sup>2</sup> area at CA-1 and AR-2 and from a 0.07-m<sup>2</sup> area at CA-2. Whole plant samples were cleaned to remove soil and other debris, and roots were separated from the stem directly above the root crown. Both plant components were dried at 60°C to a constant mass. At harvest (i.e., physiological maturity), straw and grain biomass per unit area was estimated from a 1-m<sup>2</sup> area at CA-1 and AR-2 and from a 0.07-m<sup>2</sup> area within the gas chambers at CA-2 and AR-1.

## Data Analysis

Before performing the ANOVA tests, the assumptions of normality and homogeneity of variance were evaluated. For each model, data that failed the Shapiro-Wilk test ( $p < 0.05$ ) were log transformed to meet the assumption of normality. When the data failed the Levene's test of constant variance, a weighted least squares approach was used in which the weighting factor was the reciprocal of the estimated residual variance. To determine if there were differences among cultivars in root and above-ground biomass at heading and in grain and straw weights at physiological maturity, an ANOVA was performed for each site using the PROC GLM procedure

**Table 3. Soil characteristics (0–15 cm) at the four study sites.**

Soil parameters	Study sites†			
	CA-1	CA-2	AR-1	AR-2
Taxonomic classification	Fine, smectitic, thermic Xeric Epiaquerts and Duraquerts	Fine, smectitic, thermic Typic Haploxerert	Fine, smectitic, thermic, Typic Albaqualfs	Fine, smectitic, thermic, Typic Albaqualfs
Soil series	Esquon-Neerdobe complex	Capay silty clay	Dewitt silt loam	Dewitt silt loam
Soil particle size distribution, %				
Sand	23	19	13	NM‡
Silt	30	33	69	NM
Clay	47	48	18	NM
Chemical properties				
pH	4.8	6.4	6.2	6.3
Electrical conductivity, dS m <sup>-1</sup>	0.19	NM	0.57	0.07
Total organic C, %	1.26	1.59	0.68	0.61
Total N, %	0.08	0.08	0.07	NM
Extractable P, mg kg <sup>-1</sup>	9.3	NM	13.7	8.2
Exchangeable K, mg kg <sup>-1</sup>	106	NM	198	111

† AR-1 and AR-2 refer to Arkansas field sites located at the University of Arkansas Rice Research and Extension Center near Stuttgart, AR, in 2011 and 2012, respectively; CA-1 and CA-2 refer to California field sites located at the Rice Experiment Station in Biggs in 2011 and at the University of California, Davis in 2012, respectively.

‡ Not measured.



in SAS, and unprotected Fisher's LSD was used for mean comparisons ( $p < 0.05$ ) (SAS, 2002–2010). The same analysis was used to determine if there were cultivar differences in seasonal  $\text{CH}_4$ ,  $\text{N}_2\text{O}$ , GWP, and  $\text{GWP}_Y$  within each site. For the correlation analysis, we focused on the relationships between the plant growth characteristics and seasonal  $\text{CH}_4$  emissions. We used the PROC CORR procedure in SAS to quantify Pearson's correlation coefficients ( $r$ ) of seasonal  $\text{CH}_4$  emissions of the pooled cultivars with the plant parameters at each site and used the significance levels of 0.1 and 0.05. To assess these relationships across sites, the same analysis was conducted on the pooled site data normalized to the site averages to account for differences in magnitudes of the data across sites.

## Results

### Climate

Lower seasonal air temperature and precipitation were observed at the California sites compared with the Arkansas sites. Average daily air temperatures during the rice-growing period at CA-1 and CA-2 were 21.0 and 20.4°C, respectively, compared with 26.5 and 25.2°C at AR-1 and AR-2, respectively. Cumulative seasonal precipitation rates were 0.1 and 1.4 cm at CA-1 and CA-2, respectively, and 28.8 and 40.0 cm at AR-1 and AR-2, respectively.

### Biomass Production of Cultivars

Site average grain yield across cultivars was about 32% greater on average in the California sites than in the Arkansas sites (Table 4). Average grain yields were 9.1 and 10.5  $\text{Mg ha}^{-1}$  at CA-1 and CA-2, respectively, and 6.1 and 7.2 at AR-1 and AR-2, respectively. Cultivars differed in grain yield at all three of the replicated sites (CA-1, CA-2, AR-2), but differences in straw biomass were detected at CA-1 only. At CA-1 and CA-2, M206 was the highest yielding cultivar (10.5 and 12.4  $\text{Mg ha}^{-1}$ , respectively), which was significantly greater than Calmati-202. At CA-1, M206 also had significantly greater straw biomass at maturity than L206 (9.8 and 7.9  $\text{Mg ha}^{-1}$ , respectively). At CA-2, the hybrid cultivars CLXP4534 and CLXL745 were not different from M206, the highest yielding cultivar, whereas in Arkansas, the hybrid CLXL745 was higher yielding than all the other Southern cultivars (8.3  $\text{Mg ha}^{-1}$  at AR-2).

Root and shoot biomass at heading were measured only at CA-1, CA-2, and AR-2. Site average root biomass was approximately two to three times greater at AR-2 (5.9  $\text{Mg ha}^{-1}$ ) than at CA-1 and CA-2 (2.1 and 3.0  $\text{Mg ha}^{-1}$ , respectively), whereas site average shoot biomass at heading was greatest at CA-2 (Table 4). Cultivar differences in root biomass were observed at CA-1 and CA-2 only, and these differences were not consistent. M206 and L206 had 20% greater root biomass than Calmati-202 on average at CA-1 but did not differ significantly at CA-2, although a similar trend was observed. However, at CA-2,

**Table 4. Biomass of eight cultivars at heading and physiological maturity at the four study sites.**

Cultivar	Heading		Physiological maturity	
	Root	Shoot	Grain	Straw
	Mg ha <sup>-1</sup> dry wt.			
	<b>CA-1†</b>			
Calmati-202	1.8 (0.3)‡b§	11.5 (2.2)a	7.3 (0.6)b	9.3 (0.6)ab
L206	2.3 (0.3)a	13.4 (0.9)a	9.5 (1.1)ab	7.9 (0.5)b
M206	2.2 (0.3)a	13.7 (1.4)a	10.5 (0.2)a	9.8 (0.2)a
	<b>CA-2</b>			
Calmati-202	2.5 (0.4)b	15.4 (0.8)a	7.9 (0.7)b	11.9 (0.8)a
CLXL745	3.7 (0.6)a	21.9 (2.5)a	10.9 (0.9)a	12.7 (1.2)a
CLXP4534	NM¶	NM	10.1 (0.8)ab	15.1 (3.4)a
L206	2.9 (0.1)ab	16.5 (2.8)a	10.8 (0.1)a	11.9 (0.9)a
M206	2.5 (0.2)b	16.1 (1.3)a	12.4 (1.0)a	12.8 (1.1)a
	<b>AR-1#</b>			
CLXP4534	NM	NM	7.0	11.5
Francis	NM	NM	5.4	11.7
Jupiter	NM	NM	6.9	10.1
Sabine	NM	NM	5.0	11.3
	<b>AR-2</b>			
CLXL745	6.9 (1.0)a	17.5 (3.3)a	8.3 (0.6)a	10.8 (0.6)a
Francis	4.2 (0.3)a	13.5 (0.5)ab	7.1 (0.5)b	9.8 (0.7)a
Jupiter	6.1 (0.7)a	10.8 (1.7)b	6.8 (0.7)b	10.3 (1.6)a
Sabine	6.5 (1.1)a	13.3 (1.0)ab	6.6 (0.3)b	11.2 (1.3)a

† AR-1 and AR-2 refer to Arkansas field sites located at the University of Arkansas Rice Research and Extension Center near Stuttgart, AR, in 2011 and 2012, respectively; CA-1 and CA-2 refer to California field sites located at the Rice Experiment Station in Biggs in 2011 and at the University of California, Davis in 2012, respectively.

‡ Standard errors are in parentheses.

§ Within columns of the same study site, means not sharing a letter are significantly different according to LSD (0.05).

¶ Not measured.

# Values shown for AR-1 are of a single replication and therefore no ANOVA was performed.

CLXL745 had 48% higher root biomass than Calmati-202 and M206 (Table 4). Cultivar differences in shoot biomass were detected only at AR-2; CLXL745 produced 39% more shoot biomass than Jupiter.

## N<sub>2</sub>O Emissions

Seasonal N<sub>2</sub>O emissions were negligible across all cultivars and sites (LS means of cultivars were all <0.1 kg N<sub>2</sub>O-N ha<sup>-1</sup> season<sup>-1</sup>), comprising <10% of the positive proportion of seasonal GWP when averaged across sites, and no differences were detected among cultivars within sites (Fig. 1; Table 5).

## CH<sub>4</sub> Emissions and Seasonal Patterns

Site averages at the two water-seeded sites (CA-1 and CA-2) were 63 and 13 kg CH<sub>4</sub>-C ha<sup>-1</sup>, respectively, and at the two drill-seeded sites (AR-1 and AR-2) were 34 and 70 kg CH<sub>4</sub>-C ha<sup>-1</sup>, respectively (Table 5). Seasonal patterns of daily CH<sub>4</sub> emissions were unique to each study site, and although there were observable cultivar differences in magnitudes of daily fluxes at a given site, seasonal patterns were generally similar among

cultivars at the same site, except for minor variations in timing of peak emissions (Fig. 2). In general, daily CH<sub>4</sub> fluxes started to increase after about 1 mo of flooding and began to decline toward the end of the growing season before draining. Peak CH<sub>4</sub> emissions during the flooded period generally occurred in early August for all cultivars across all sites after about 2.5 mo of continuous flooding, except at AR-1, where the first peak in CH<sub>4</sub> emissions occurred after 1.5 mo of flooding. Flooding was delayed (day of year) at AR-1 compared with all other sites, although the interval between seeding and flooding at AR-1 (34 d) was shorter than at AR-2 (36 d). Furthermore, we observed two peaks at AR-1, the first 1.5 mo after flooding in early August and the second, similar to the other sites, about 2 mo after flooding in late August. After a sharp decline in CH<sub>4</sub> flux from 18 to 23 August at CA-2, a series of attenuated CH<sub>4</sub> fluxes was observed. After drainage at the end of the rice-growing season, spikes in CH<sub>4</sub> emissions were measured at all sites. However, the magnitudes of the peaks at CA-2 and AR-2 were relatively minor compared with the magnitude of the fluxes preceding the drainage event.




Fig. 1. Seasonal patterns of daily N<sub>2</sub>O emissions of eight cultivars and air temperature during the rice-growing season at CA-1, CA-2, AR-1, and AR-2 (AR-1 and AR-2 refer to Arkansas field sites located at the University of Arkansas Rice Research and Extension Center near Stuttgart, AR, in 2011 and 2012, respectively; CA-1 and CA-2 refer to California field sites located at the Rice Experiment Station in Biggs in 2011 and at the University of California, Davis in 2012, respectively). Data are plotted on different scales for each plot. Error bars represent SE of the replicates. If no error bar is indicated, only one replicate was available. Daily mean air temperature is plotted above the emissions profile in each graph panel. Dates of seeding (arrow), N fertilization (F), the continuously flooded period (shaded), and harvest (M) are shown. Heading dates (H) of the cultivars are indicated as follows: (CA-1) H = all cultivars (15–21 Aug.), (CA-2) H = all cultivars (4–12 Sept.), (AR-1) H1 = CLXP4534 (31 July), H2 = all other cultivars (10–12 Aug.), (AR-2) H1 = CLXL745 (11–14 July), and (AR-2) H2 = all other cultivars (15–20 July).

There was no clear relationship between heading of the rice plants and seasonal patterns in CH<sub>4</sub> emissions (Fig. 2). Whereas the relationship between the heading of individual cultivars and peak CH<sub>4</sub> emissions during the flooded period depended on the site and cultivar, the difference in magnitude of cumulative emissions before and after heading depended on the site (Fig. 2; Table 5). During the flooded period at all sites, heading for each cultivar occurred at the beginning of, or within 1 mo after, the maximum CH<sub>4</sub> flux except at AR-2 where heading occurred within 2 wk before peak CH<sub>4</sub> emissions for all cultivars. Regardless of cultivar, in the CA water-seeded systems the majority of seasonal CH<sub>4</sub> emissions occurred before the heading stage (>76% on average) (Table 5). Conversely, in the AR drill-seeded systems, all the cultivars except Jupiter at AR-1 had a higher proportion of seasonal CH<sub>4</sub> emissions occur after heading (>72% on average) (Table 5).

### Cultivar Comparisons in Seasonal CH<sub>4</sub> within Sites

The greatest range in seasonal emissions among cultivars at the same site was observed at AR-1, with 15 kg CH<sub>4</sub>-C ha<sup>-1</sup> in Jupiter and 55 kg CH<sub>4</sub>-C ha<sup>-1</sup> in Sabine (Table 5). However, at AR-1 these values represent a single replication. At AR-2 there was only a 3% difference in mean seasonal CH<sub>4</sub> emissions of these same cultivars; this difference was not statistically significant. At the water-seeded sites (CA-1 and CA-2), Calmati-202 tended to have the lowest mean seasonal CH<sub>4</sub> emissions, but it was

only significantly lower than L206 at one of the sites (CA-2). The hybrid cultivars CLXP4534 and CLXL745 were moderate emitters of CH<sub>4</sub> at CA-2 relative to the other cultivars, whereas CLXL745 had the lowest seasonal CH<sub>4</sub> emissions at AR-2 (56 kg CH<sub>4</sub>-C ha<sup>-1</sup>) ( $p < 0.05$ ), and CLXP4534 had the second to lowest at AR-1 (25 kg CH<sub>4</sub>-C ha<sup>-1</sup>). At AR-2 the disparity between CLXL745 and the other cultivars in CH<sub>4</sub> emissions occurred at the end of the season, starting in late July (Fig. 2).

### Global Warming Potential

Seasonal GWP was strongly related to CH<sub>4</sub> emissions at all sites, but N<sub>2</sub>O emissions were only a minor contributor to GWP (Table 5). Site average seasonal GWP values at the water-seeded sites (CA-1 and CA-2) were 2144 and 314 kg CO<sub>2</sub> eq ha<sup>-1</sup>, respectively, and at the drill-seeded sites (AR-1 and AR-2) were 1254 and 2399 kg CO<sub>2</sub> eq ha<sup>-1</sup>, respectively. At least 90% of seasonal GWP at each site on average was due to CH<sub>4</sub> emissions; this finding was consistent among cultivars.

Seasonal GWP<sub>Y</sub> was similar to GWP and CH<sub>4</sub> emissions in terms of relative differences among sites and cultivars. The CA-2 had the lowest mean GWP<sub>Y</sub> (24 kg CO<sub>2</sub>-eq Mg<sup>-1</sup>), which was about 14 times less than the site average at AR-2 (339 kg CO<sub>2</sub>-eq Mg<sup>-1</sup>) (Table 5). CA-1 and AR-1 had similar mean GWP<sub>Y</sub> (240 and 222 kg CO<sub>2</sub>-eq Mg<sup>-1</sup>, respectively). The greatest variability in GWP<sub>Y</sub> among cultivars was observed at CA-2, with an approximately 3-fold difference in GWP<sub>Y</sub> between L206 and

**Table 5. Seasonal CH<sub>4</sub> and N<sub>2</sub>O fluxes, global warming potential, and yield-scaled global warming potential of eight cultivars at the four study sites.**

Cultivar	Seasonal CH <sub>4</sub> kg CH <sub>4</sub> -C ha <sup>-1</sup>	Seasonal N <sub>2</sub> O kg N <sub>2</sub> O-N ha <sup>-1</sup>	Seasonal GWP† kg CO <sub>2</sub> eq ha <sup>-1</sup>	Seasonal GWP‡ kg CO <sub>2</sub> eq <sup>-1</sup> Mg <sup>-1</sup>	Heading date(s)	Cumulative CH <sub>4</sub> emissions	
						Before heading	After heading
— kg CH <sub>4</sub> -C ha <sup>-1</sup> —							
<b>CA-1§</b>							
Calmati-202	58 (2.9)a¶	0.05 (0.021)a	1988 (90)a	277 (19)a	20 Aug.	32 (4.5)a	26 (2.9)a
L206	60 (12)a	0.06 (0.004)a	2074 (399)a	218 (29)a	21 Aug.	42 (12)a	19 (2.9)b
M206	69 (9.6)a	0.06 (0.018)a	2370 (319)a	225 (30)a	15 Aug.	44 (10)a	25 (0.73)a
<b>CA-2</b>							
Calmati-202	6.7 (0.9)b	-0.15 (0.016)a	83 (31)b	10 (3.6)b	12 Sept.	5.9 (0.7)b	0.80 (0.2)b
CLXL745	11 (3.1)ab	-0.19 (0.025)a	309‡ (99)ab	29# (12)ab	4 Sept.	8.5 (2.3)ab	2.5 (0.8)a
CLXP4534	11 (0.4)ab	-0.16 (0.037)a	229 (40)ab	22 (2.6)ab	12 Sept.	10 (0.4)a	1.2 (0.01)ab
L206	14 (2.2)a	-0.13 (0.016)a	353 (62)a	33 (6.1)a	4 Sept.	11 (1.1)a	3.5 (1.3)a
M206	11 (1.1)ab	-0.11 (0.028)a	277 (61)ab	23 (6.1)ab	4 Sept.	8.6 (0.2)ab	2.8 (0.9)a
<b>AR-1††</b>							
CLXP4534	25	0.17	981	140	31 July	4.2	20
Francis	43	0.10	1526	282	10 Aug.	16	27
Jupiter	15	0.08	573	83	12 Aug.	7.7	7.2
Sabine	55	0.10	1936	384	12 Aug.	24	31
<b>AR-2</b>							
CLXL745	56 (9.2)b	0.02 (0.047)a	1899 (344)b	232 (44)b	11–14 July	16 (4.9)a	40 (4.4)b
Francis	77 (14)a	0.10 (0.052)a	2677 (459)a	381 (81)a	15–20 July	18 (4.0)a	59 (10)a
Jupiter	72 (16)ab	-0.01 (0.022)a	2397 (534)ab	345 (59)ab	20–22 July	29 (8.8)a	43 (7.0)b
Sabine	75 (7.5)ab	0.11 (0.114)a	2623 (154)a	397 (30)a	19–20 July	37 (11)a	38 (11)b

† Global warming potential.


‡ Yield-scaled global warming potential.

§ AR-1 and AR-2 refer to Arkansas field sites located at the University of Arkansas Rice Research and Extension Center near Stuttgart, AR, in 2011 and 2012, respectively; CA-1 and CA-2 refer to California field sites located at the Rice Experiment Station in Biggs in 2011 and at the University of California, Davis in 2012, respectively.

¶ Within columns of the same study site, means not sharing a letter are significantly different according to LSD (0.05). Standard errors are in parentheses.

# Mean of two replicates due to removal of outlier.

†† Values shown for AR-1 are of a single replicate, and therefore no ANOVA was performed.



**Fig. 2.** Seasonal patterns of daily  $\text{CH}_4$  emissions of eight cultivars and air temperature during the rice-growing season at CA-1, CA-2, AR-1, and AR-2 (AR-1 and AR-2 refer to Arkansas field sites located at the University of Arkansas Rice Research and Extension Center near Stuttgart, AR, in 2011 and 2012, respectively; CA-1 and CA-2 refer to California field sites located at the Rice Experiment Station in Biggs in 2011 and at the University of California, Davis in 2012, respectively). The data are plotted on different scales for each plot. Error bars represent the SE of the replicates. If no error bar is indicated, only one replicate was available. Daily mean air temperature is plotted above the emissions profile in each graph panel. Dates of seeding (arrow), N fertilization (F), the continuously flooded period (shaded), and harvest (M) are shown. Heading dates (H) of the cultivars are indicated as follows: (CA-1) H = all cultivars (15–21 Aug.), (CA-2) H = all cultivars (4–12 Sept.), (AR-1) H1 = CLXP4534 (31 July), H2 = all other cultivars (10–12 Aug.), (AR-2) H1 = CLXL745 (11–14 July), and (AR-2) H2 = all other cultivars (15–20 July).

Calmati-202 (32.7 and 10.0 kg  $\text{CO}_2\text{-eq Mg}^{-1}$ , respectively). Variability in  $\text{GWP}_Y$  among the cultivars at CA-1 was minimal, ranging from 218 to 277 kg  $\text{CO}_2\text{-eq Mg}^{-1}$ , and no differences were detected among cultivars. At AR-2, the hybrid CLXL745 had the lowest  $\text{GWP}_Y$  (232 kg  $\text{CO}_2\text{-eq}^{-1} \text{Mg}^{-1}$ ), which was significantly less than Sabine and Francis (396.7 and 380.7 kg  $\text{CO}_2\text{-eq}^{-1} \text{Mg}^{-1}$ , respectively).

### Correlations between Plant Growth Parameters and Seasonal $\text{CH}_4$ Emissions

The relationships between seasonal  $\text{CH}_4$  emissions and the plant parameters of the pooled cultivars were not consistent across sites (Table 6); seasonal  $\text{CH}_4$  emissions were not significantly correlated with any plant parameters at CA-2 or AR-2 but were positively correlated with shoot and total plant (root + shoot) biomass at heading at CA-1 ( $r = 0.60$ ) and negatively correlated with grain yield at AR-1 ( $r = -0.95$ ). Excluding AR-1 (where no root measurements were taken), there was a positive correlation ( $r = 0.33$ ) between root biomass at heading and seasonal  $\text{CH}_4$

emissions when sites were pooled and data were normalized to the corresponding site averages.

## Discussion

### Relative Importance of $\text{CH}_4$ and $\text{N}_2\text{O}$ to Global Warming Potential

Our research shows that  $\text{CH}_4$  emissions contributed at least 90% to seasonal GWP when cultivars were averaged within sites, which is consistent with other field studies on GHG emissions in wet- and drill-seeded rice systems (Adviento-Borbe et al., 2013; Pittelkow et al., 2013) and with the findings from a meta-analysis of 16 studies in transplanted rice systems (Linguist et al., 2012). Seasonal  $\text{N}_2\text{O}$  emissions included negative and positive fluxes but were minimal (Table 5). Consumption of  $\text{N}_2\text{O}$  in anoxic soils is via denitrification when  $\text{N}_2\text{O}$  is reduced to  $\text{N}_2$ . Such fluxes from the atmosphere to soil have been reported from rice systems (Chen et al., 1997; Xu et al., 1997, 2004) and from natural ecosystems, with the highest values from wetland and peatland ecosystems (Schlesinger, 2013). Our estimate of  $\text{N}_2\text{O}$  contribution to GWP may be underestimated at the AR sites due



to gas sampling not starting until after the onset of the permanent flood, approximately 6 wk after planting and 2 to 3 wk after N fertilization, which likely missed some peaks in N<sub>2</sub>O emissions. For example, at CA-2 the highest peak in N<sub>2</sub>O emissions was observed approximately 2 wk after N fertilization. However, this flux was observed on the same day as water seeding, which may have been an artifact of that disturbance. Indeed, other studies in drill- and water-seeded systems have found low to undetectable N<sub>2</sub>O emissions after N fertilization when fields are immediately flooded (Adviento-Borbe et al., 2013; Pittelkow et al., 2013), which is to be expected from flooded soils due to their low redox potential (Hou et al., 2000). Nitrous oxide emissions are typically low before flooding because the fertilizer N has not yet been applied (Adviento-Borbe et al., 2013). Nitrous oxide emissions during this period likely played a relatively insignificant role in the cultivar differences in GWP because without surface floodwater the primary pathway for N<sub>2</sub>O emissions is through the soil (Yan et al., 2000) and not through the plant.

Because the objective of this study was to evaluate cultivar differences in seasonal GHG emissions, postharvest N<sub>2</sub>O and CH<sub>4</sub> emissions were not accounted for. The annual effect of cultivar on GWP is likely determined by emissions during the growing season when the plants are present. Although N<sub>2</sub>O fluxes after harvest may have persisted depending on soil moisture status, precipitation, and N availability, we would not expect the additional emissions to greatly influence total cumulative seasonal N<sub>2</sub>O emissions or GWP (CH<sub>4</sub> + N<sub>2</sub>O) of cultivars.

### Cumulative CH<sub>4</sub> Emissions and Seasonal Patterns among Sites

The site average seasonal CH<sub>4</sub> emissions at three of the four sites (CA-1, AR-1, AR-2) ranged from 34 to 70 kg CH<sub>4</sub>-C ha<sup>-1</sup> season<sup>-1</sup>, which is within the lower ranges reported for U.S. rice systems (Bossio et al., 1999; Fitzgerald et al., 2000; Sass et al., 2002; Pittelkow et al., 2013) (Table 5). Seasonal CH<sub>4</sub> emissions at CA-2, on the other hand, were particularly low (site average of 13 kg CH<sub>4</sub>-C ha<sup>-1</sup>), likely because this field had been fallow the previous 4 yr and residue inputs into this system were limited to weeds only, reducing C substrate and ultimately lowering CH<sub>4</sub> production potential in this soil (Zhanget al., 2013). Additionally,

the field was not flooded during the extended fallow period. Low historic inputs of organic matter and lack of previous water-logged conditions have been shown to reduce CH<sub>4</sub> emissions in subsequent rice-growing seasons (Yan et al., 2005). Furthermore, although CA-2 had higher soil organic C (SOC) content than all other sites, the clay content was 48%, which is known to adsorb and protect SOC from decomposition (Baldock and Skjemstad, 2000), further decreasing the methanogenic source strength of this soil. High clay content is also known to increase entrapment of CH<sub>4</sub> (Wang et al., 1993), ultimately lowering CH<sub>4</sub> transport from the soil to the atmosphere. Although there is less than a 1°C difference in mean seasonal daily air temperature between the two CA sites, CA-2 was 6.1 and 4.8°C lower than AR-1 and AR-2, respectively, perhaps also contributing to the relatively low CH<sub>4</sub> emissions.

The seasonal trends in daily CH<sub>4</sub> emissions do not appear to be driven by development stage of the rice plant because heading was observed to occur before, during, and after peaks in CH<sub>4</sub> emissions (Fig. 2). Trends in daily CH<sub>4</sub> emissions also do not appear to consistently respond to changes in air temperature (Fig. 2). The dual CH<sub>4</sub> peaks observed at CA-2 and AR-1 can also be seen in the seasonal CH<sub>4</sub> patterns of other studies (Chidthaisong and Watanabe, 1997; Wassmann et al., 2002; Pittelkow et al., 2014). In a <sup>13</sup>C-tracer experiment, Chidthaisong and Watanabe (1997) observed two CH<sub>4</sub> peaks. The first peak occurred 40 to 80 d after flooding and was related to the previous seasons' straw residue-derived CH<sub>4</sub> but was primarily derived from SOC and the roots. A second peak occurred past 80 d after flooding, at the heading stage, and represented the increase in emissions driven by an increase in plant-derived C sources with relatively minimal contribution from straw and SOC. This suggests that the C source of initial CH<sub>4</sub> emissions was from all C pools, including roots, SOC, and plant residue, whereas for late-season emissions the C source was primarily from root-derived organic material.

All sites showed a decline in CH<sub>4</sub> emissions before draining, which could be due to a number of factors related to increased CH<sub>4</sub> oxidation, decreased CH<sub>4</sub> production, and decreased CH<sub>4</sub> transport capacity of the rice plants (Fig. 2). The proportion of CH<sub>4</sub> oxidized after heading (30–70%) increased compared with pre-heading (10–20%) (Sigren et al., 1997). In a greenhouse

**Table 6.** Pearson's correlation coefficients of the pooled cultivar seasonal CH<sub>4</sub> emissions with plant biomass parameters within and across study sites.

Study site†	Sample size (n)	Heading			Physiological maturity		
		Root	Shoot	Total (root + shoot)	Grain	Straw	Total (grain + straw)
CA-1	9	0.45	0.60‡	0.60‡	0.53	0.27	0.51
CA-2	12§, 15¶	0.40	0.12	0.17	0.37	0.08	0.28
AR-1	4	NM#	NM	NM	−0.95*	0.63	−0.58
AR-2	12	0.20	−0.47	−0.35	−0.14	0.17	0.07
All sites††	33§, 40¶	0.33‡	−0.03	0.09	0.09	0.14	0.23

\* Significant at the 0.05 probability level.

† AR-1 and AR-2 refer to Arkansas field sites located at the University of Arkansas Rice Research and Extension Center near Stuttgart, AR, in 2011 and 2012, respectively; CA-1 and CA-2 refer to California field sites located at the Rice Experiment Station in Biggs in 2011 and at the University of California, Davis in 2012, respectively.

‡ Significant at the 0.1 probability level.

§ For root or shoot.

¶ For grain and straw.

# Not measured; root and shoot measurements were not taken at AR-1.

†† All data normalized to the corresponding site mean.

study on the CH<sub>4</sub> transport capacity of rice plants, Aulakh et al. (2000) reported a decline in CH<sub>4</sub> transport capacity from flowering to maturity of all four high-yielding dwarf cultivars (IR72, IR52, IR64, and PSBRc20), two of the three traditional tall cultivars (Dular and B40), and one of the three new plant type cultivars (IR65598).

## Cultivar Differences in CH<sub>4</sub> emissions, Global Warming Potential, and Yield-Scaled Global Warming Potential

Methane emissions from flooded rice soils are the net effect of plant-regulated mechanisms that affect CH<sub>4</sub> production, oxidation, and transport of gases between the rhizosphere and atmosphere and their interaction with the environment and management factors. Some studies have found no persistent cultivar effect on CH<sub>4</sub> emissions due to strong interactions with the environment (e.g., Wassmann et al., 2002). Our study of two site-years in Arkansas and California provides support that relative differences or similarities in seasonal CH<sub>4</sub> among some cultivars are consistent across sites, whereas other cultivars are not. At both CA sites, M206 and L206 were similar, whereas Calmati-202 tended to be lower but only significantly lower than L206 at one site (CA-2). Although the statistical significance at the AR sites could only be tested at the replicated site (AR-2), a similar trend was observed among the Southern cultivars at both sites; Francis and Sabine were similar in seasonal CH<sub>4</sub> emissions and were greater than the hybrid (CLXP4534 or CLXL745). In contrast, Jupiter was more variable in seasonal CH<sub>4</sub> emissions relative to the other cultivars at AR-1 and AR-2, and the ranking of this cultivar in comparison to the hybrid was reversed across sites. All of the cultivars studied are being commercially produced and represent a very narrow and elite subset of rice germplasm that is available. Although few statistical differences among the cultivars were observed, this may not reflect the genetic diversity that is available for mitigating GHG emissions. A better understanding of the genotype × environment interaction is necessary to determine to what extent choice of rice cultivar may affect GWP relative to cultural management and environmental factors.

Watanabe et al. (1995) did not find any correlation between CH<sub>4</sub> emissions and shoot or root biomass despite significant cultivar differences in CH<sub>4</sub> emissions. However, in our study there was a small positive correlation of root biomass at heading with seasonal CH<sub>4</sub> emissions when evaluated across sites (Table 6). This relationship was similar to findings of Wang et al. (1997), who attributed it to total C released from roots, indicating that the cultivar differences in CH<sub>4</sub> emissions were mainly due to differences in their effect on CH<sub>4</sub> production.

In a study investigating microbial controls on differences in CH<sub>4</sub> emissions among three rice cultivars, Ma et al. (2010) found that the cultivar with higher shoot biomass had lower CH<sub>4</sub> emissions due to its higher CH<sub>4</sub> oxidation potential. In contrast, our research shows a positive correlation between seasonal CH<sub>4</sub> emissions and shoot biomass at CA-1, although there were no differences in shoot biomass or CH<sub>4</sub> emissions among cultivars at CA-1 (Tables 4 and 5). Variability in the mechanisms driving CH<sub>4</sub> emissions, and in the associated plant traits, may be just as great within cultivars as the variability among cultivars due to genotypic plasticity in response to micro-environment effects.

On the other hand, at CA-2 and AR-2, cultivars differed in seasonal CH<sub>4</sub> emissions (Table 5) and in root and shoot biomass, respectively (Table 4), but there were no significant correlations between CH<sub>4</sub> emissions and the measured plant parameters at either site (Table 6). These findings highlight that plant biomass parameters alone are not dependable proxies for relative differences in CH<sub>4</sub> emissions or the processes driving CH<sub>4</sub> emissions.

Among cultivars with similar root and shoot biomass at heading, we expected those with higher grain yield to have lower CH<sub>4</sub> emissions based on the reasoning that CH<sub>4</sub> transport would be similar but CH<sub>4</sub> production would be less in the higher yielding cultivar due to greater allocation of C to grain versus root exudation. Denier van der Gon et al. (2002) observed an increase in CH<sub>4</sub> emissions after removing spikelets and thereby artificially decreasing yield potential by lowering the capacity of the reproductive organs to accept C assimilates. In other words, the yield gap (the expected grain yield based on the maximum estimated harvest index times the total above-ground biomass minus actual yield) was proportional to CH<sub>4</sub> emissions, which is presumably via leakage of excess photosynthetically fixed C from roots. This may explain the significant negative correlation ( $r = -0.96$ ) between CH<sub>4</sub> emissions and yield at AR-1 (Table 6). Data supporting this hypothesis were observed at AR-2, in which CLXL745 had similar root and shoot biomass, higher yield, and the lowest CH<sub>4</sub> emissions compared with Francis and Sabine. The disparity between CLXL745 and the other cultivars occurred at the end of the season (Fig. 2) when root-derived C has been shown to be the main contributor to CH<sub>4</sub> emissions (Chidthaisong and Watanabe, 1997). However, a general conclusion on the net effect of yield on seasonal CH<sub>4</sub> emissions in cultivars with similar root and shoot biomass cannot be made due to conflicting results at CA-2, in which L206 had similar root and shoot biomass, higher yield, and higher CH<sub>4</sub> emissions compared with Calmati-202 (Tables 4 and 5).

Evaluating differences in GHG emissions from agriculture relative to yield (GWP<sub>y</sub>), as opposed to land area alone (GWP), has become increasingly relevant due to the recognition that global food security and environmental health should be considered in evaluating GHG mitigation options (van Groenigen et al., 2010; Grassini and Cassman, 2012; Linquist et al., 2012). The optimal scenario for a cultivar with the lowest GWP<sub>y</sub> is for it to have the highest yield and the lowest GWP and for these characteristics to persist across environments and years. Three Southern hybrid cultivars, including CLXL745 tested in the present study, have been shown to yield 1 to 1.7 Mg ha<sup>-1</sup> more than the highest-yielding conventional alternative (Francis) in Arkansas (Lyman and Nalley, 2013), which is corroborated by our results at AR-1 and AR-2 (Table 4). Thus, given that the lowest CH<sub>4</sub>-emitting cultivar at AR-2 was the hybrid, CLXL745, and that hybrids tend to yield more than conventional Southern cultivars, lower GWP<sub>y</sub> values are expected (Table 5). This shows there is potential for breeding high-yielding cultivars with low GWP and GWP<sub>y</sub>.

## Conclusions

Development of carbon offset protocols for GHG mitigation in agriculture is underway, such as for rice systems in California (Climate Action Reserve, 2013). Process-based biogeochemical models are proposed for estimation of GHG emissions. There

are options in some models to customize plant parameters (e.g., C:N ratios of biomass, biomass ratios, and maximum yield), but the variation in the mechanisms driving cultivar differences in GHG emissions (i.e., CH<sub>4</sub> production, oxidation, and transport capacity) are not explicit. We found that, within study sites, cultivars were similar in seasonal N<sub>2</sub>O emissions due to the continuously flooded conditions for the majority of plant growth and development, and most cultivars were also similar in seasonal CH<sub>4</sub> emissions. However, there were two cases in which cultivars differed in seasonal CH<sub>4</sub> emissions by a factor of 2.1 and 1.4. Accordingly, cultivar differences in GWP were similar to differences in CH<sub>4</sub> emissions and not N<sub>2</sub>O emissions. Our research shows that relative seasonal CH<sub>4</sub> emissions among some cultivars were consistent across two site-years, whereas other cultivars varied, presumably due to varying degrees to which cultivar phenotypic characteristics related to CH<sub>4</sub> emissions respond to external factors. We found contradictory relationships between combinations of crop growth characteristics and seasonal CH<sub>4</sub> emissions, highlighting that plant biomass parameters alone are not dependable proxies for relative differences in CH<sub>4</sub> emissions or for the processes driving CH<sub>4</sub> emissions due to complex interactions among plant phenotypes, the environment, and mechanisms driving CH<sub>4</sub> production, oxidation, and transport. At one of the Arkansas sites, the hybrid had the highest yield, similar root and shoot biomass at heading, and lower GWP and GWP<sub>Y</sub> compared with two of the three inbred cultivars. This result may have important implications for breeding high-yielding cultivars with low GWP and GWP<sub>Y</sub>.

## Acknowledgments

The research was funded in part by Mars, Inc., and by the William G. and Kathleen Golden International Agriculture Fellowship, UCD & Humanities Graduate Research Award, and the Department of Plant Sciences at the Univ. of California, Davis, through a graduate student research assistantship awarded to Maegen Simmonds. We thank the Agroecosystems Lab, especially Cesar Abrenilla, Christopher Mikita, Lijun Liu, Xinqiang Liang, Toni Leong, Alec Liou, Ian Marton, Evonne Soon, Emilie Kirk, and Gabriel La Hue, as well as Shelly Kerr and David Hendrix from the Univ. of Arkansas; employees at the Dale Bumpers National Rice Research Center; and Ray Stogsdill at the California Rice Experiment Station, for their tremendous support in the lab and/or field. We greatly appreciate Cameron Pittelkow, Yacov Assa, and Tim Doane for their technical assistance and Johan Six for sharing equipment. We also greatly appreciate the statistical consultations provided by Neil Willits from the Univ. of California, Davis, and Kathleen Yeater from the USDA-ARS.

## References

Adviento-Borbe, M.A., C.M. Pittelkow, M. Anders, C. van Kessel, J.E. Hill, A.M. McClung, J. Six, and B.A. Linquist. 2013. Optimal fertilizer nitrogen rates and yield-scaled global warming potential in drill seeded rice. *J. Environ. Qual.* 42:1623–1634. doi:10.2134/jeq2013.05.0167

Aulakh, M.S., J. Bodenbender, R. Wassmann, and H. Rennenberg. 2000. Methane transport capacity of rice plants: II. Variations among different rice cultivars and relationship with morphological characteristics. *Nutr. Cycling Agroecosyst.* 58:367–375. doi:10.1023/A:1009839929441

Aulakh, M.S., R. Wassmann, C. Bueno, J. Kreuzwieser, and H. Rennenberg. 2001a. Characterization of root exudates at different growth stages of ten rice (*Oryza sativa* L.) cultivars. *Plant Biol.* 3:139–148. doi:10.1055/s-2001-12905

Aulakh, M.S., R. Wassmann, C. Bueno, and H. Rennenberg. 2001b. Impact of root exudates of different cultivars and plant developmental stages of rice (*Oryza sativa* L.) on methane production in a paddy soil. *Plant Soil* 230:77–86. doi:10.1023/A:1004817212321

Baldock, J.A., and J.O. Skjemstad. 2000. Role of the soil matrix and minerals in protecting natural organic materials against biological attack. *Org. Geochem.* 31:697–710. doi:10.1016/S0146-6380(00)00049-8

Bilek, R., S.C. Tyler, R.L. Sass, and F.M. Fisher. 1999. Differences in CH<sub>4</sub> oxidation and pathways of production between rice cultivars deduced from measurements of CH<sub>4</sub> flux and δ<sup>13</sup>C of CH<sub>4</sub> and CO<sub>2</sub>. *Global Biogeochem. Cycles* 13:1029–1044. doi:10.1029/1999GB900040

Bossio, D.A., W.R. Horwath, R.G. Mutters, and C. van Kessel. 1999. Methane pool and flux dynamics in a rice field following straw incorporation. *Soil Biol. Biochem.* 31:1313–1322. doi:10.1016/S0038-0717(99)00050-4

Butterbach-Bahl, K., H. Papen, and H. Rennenberg. 1997. Impact of gas transport through rice cultivars on methane emission from rice paddy fields. *Plant Cell Environ.* 20:1175–1183. doi:10.1046/j.1365-3040.1997.d01-142.x

Chen, G.X., G.H. Huang, B. Huang, K.W. Yu, J. Wu, and H. Xu. 1997. Nitrous oxide and methane emissions from soil-plant systems. *Nutr. Cycling Agroecosyst.* 49:41–45. doi:10.1023/A:1009758900629

Chen, Y.-H., and R.G. Prinn. 2006. Estimation of atmospheric methane emission between 1996 and 2001 using a three-dimensional global chemical transport model. *J. Geophys. Res.* 111:D10307. doi:10.1029/2005JD006058

Chidthaisong, A., and I. Watanabe. 1997. Methane formation and emission from flooded rice soil incorporated with <sup>13</sup>C-labeled rice straw. *Soil Biol. Biochem.* 29:1173–1181. doi:10.1016/S0038-0717(97)00034-5

Climate Action Reserve. 2013. Rice cultivation project protocol. Version 1.1. Climate Action Reserve, Los Angeles, CA. <http://www.climateactionreserve.org/how/protocols/rice-cultivation/> (13 Mar. 2014).

Colmer, T.D. 2003. Long-distance transport of gases in plants: A perspective on internal aeration and radial oxygen loss from roots. *Plant Cell Environ.* 26:17–36. doi:10.1046/j.1365-3040.2003.00846.x

Conrad, R. 2005. Quantification of methanogenic pathways using stable carbon isotopic signatures: A review and a proposal. *Org. Geochem.* 36:739–752. doi:10.1016/j.orggeochem.2004.09.006

Denier van der Gon, H.A.C., M.J. Kropff, N. van Breemen, R. Wassmann, R.S. Lantin, E. Aduna, T.M. Corton, and H.H. van Laar. 2002. Optimizing grain yields reduces CH<sub>4</sub> emissions from rice paddy fields. *Proc. Natl. Acad. Sci. USA* 99:12021–12024. doi:10.1073/pnas.192276599

Eller, G., and P. Frenzel. 2001. Changes in activity and community structure of methane-oxidizing bacteria over the growth period of rice. *Appl. Environ. Microbiol.* 67:2395–2403. doi:10.1128/AEM.67.6.2395-2403.2001

Fitzgerald, G.J., K.M. Scow, and J.E. Hill. 2000. Fallow season straw and water management effects on methane emissions in California rice. *Global Biogeochem. Cycles* 14:767–776. doi:10.1029/2000GB001259

Forster, P., V. Ramaswamy, P. Artaxo, T. Berntsen, R. Betts, D.W. Fahey, J. Haywood, J. Lean, D.C. Lowe, G. Myhre, J. Nganga, R. Prinn, G. Raga, M. Schulz, and R. Van Dorland. 2007. 2007: Changes in atmospheric constituents and in radiative forcing. In: S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor, and H.L. Miller, editors, *Climate change 2007: The physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge Univ. Press, Cambridge, UK, and New York, p. 129–234.

Grassini, P., and K.G. Cassman. 2012. High-yield maize with large net energy yield and small global warming intensity. *Proc. Natl. Acad. Sci. USA* 109:1074–1079. doi:10.1073/pnas.1116364109

Holzappel-Pschorn, A., R. Conrad, and W. Seiler. 1986. Effects of vegetation on the emission of methane from submerged paddy soil. *Plant Soil* 92:223–233. doi:10.1007/BF02372636

Holzappel-Pschorn, A., and W. Seiler. 1986. Methane emission during a cultivation period from an Italian rice paddy. *J. Geophys. Res.* 91:11803–11814. doi:10.1029/JD091iD11p11803

Hou, A.X., G.X. Chen, Z.P. Wang, O. Van Cleemput, and W.H. Patrick, Jr. 2000. Methane and nitrous oxide emissions from a rice field in relation to soil redox and microbiological processes. *Soil Sci. Soc. Am. J.* 64:2180–2186. doi:10.2136/sssaj2000.6462180x

Huang, Y., R.L. Sass, and F.M. Fisher, Jr. 1997. Methane emission from Texas rice paddy soil: I. Quantitative multi-year dependence of CH<sub>4</sub> emission on soil, cultivar and grain yield. *Glob. Change Biol.* 3:479–489. doi:10.1046/j.1365-2486.1997.00083.x

Hutchinson, G.L., and G.P. Livingston. 1993. Use of chamber systems to measure trace gas fluxes. In: D.E. Rolston, J.M. Duxbury, L.A. Harper, and R. Mosier, editors, *Agricultural ecosystem effects on trace gases and global climate change*. ASA Spec. Publ. 55. ASA, CSSA, SSSA, Madison, WI, p. 63–78.

Linquist, B.A., K.J. van Groenigen, M.A. Adviento-Borbe, C. Pittelkow, and C. van Kessel. 2012. An agronomic assessment of greenhouse gas emissions from major cereal crops. *Glob. Change Biol.* 18:194–209. doi:10.1111/j.1365-2486.2011.02502.x



- Lu, W.F., W. Chen, B.W. Duan, W.N. Guo, Y. Lu, R.S. Lantin, R. Wassmann, and H.U. Neue. 2000a. Methane emissions and mitigation options in irrigated rice fields in southeast China. *Nutr. Cycling Agroecosyst.* 58:65–73. doi:10.1023/A:1009830232650
- Lu, Y., R. Wassmann, H.-U. Neue, and C. Huang. 2000b. Dynamics of dissolved organic carbon and methane emissions in a flooded rice soil. *Soil Sci. Soc. Am. J.* 64:2011–2017. doi:10.2136/sssaj2000.6462011x
- Lu, Y., A. Watanabe, and M. Kimura. 2002. Contribution of plant-derived carbon to soil microbial biomass dynamics in a paddy rice microcosm. *Biol. Fertil. Soils* 36:136–142. doi:10.1007/s00374-002-0504-2
- Lyman, N., and L.L. Nalley. 2013. Economic analysis of hybrid rice performance in Arkansas. *Agron. J.* 105:977–988. doi:10.2134/agronj2012.0461
- Ma, K., Q. Qiu, and Y. Lu. 2010. Microbial mechanism for rice control on methane emission from rice field soil. *Glob. Change Biol.* 16:3085–3095. doi:10.1111/j.1365-2486.2009.02145.x
- Parkin, T.B., R.T. Venterea, and S.K. Hargreaves. 2012. Calculating the detection limits of chamber-based soil greenhouse gas flux measurements. *J. Environ. Qual.* 41:705–715. doi:10.2134/jeq2011.0394
- Pittelkow, C.M., M.A. Adviento-Borbe, J.E. Hill, J. Six, C. van Kessel, and B.A. Linquist. 2013. Yield-scaled global warming potential of annual nitrous oxide and methane emissions from continuously flooded rice in response to nitrogen input. *Agric. Ecosyst. Environ.* 177:10–20. doi:10.1016/j.agee.2013.05.011
- Pittelkow, C.M., Y. Assa, M. Burger, R.G. Mutters, C.A. Greer, L.A. Espino, J.E. Hill, W.R. Horwath, C. van Kessel, and B.A. Linquist. 2014. Nitrogen management and methane emissions in direct-seeded rice systems. *Agron. J.* 106:968–980. doi:10.2134/agronj13.0491
- SAS. 2002–2010. SAS System for Windows, Version 9.3. SAS Inst., Inc., Cary, NC.
- Sass, R.L., J.A. Andrews, A. Ding, and F.M. Fisher, Jr. 2002. Spatial and temporal variability in methane emissions from rice paddies: Implications for assessing regional methane budgets. *Nutr. Cycling Agroecosyst.* 64:3–7. doi:10.1023/A:1021107016714
- Schlesinger, W.H. 2013. An estimate of the global sink for nitrous oxide in soils. *Glob. Change Biol.* 19:2929–2931. doi:10.1111/gcb.12239
- Sigren, L.K., G.T. Byrd, F.M. Fisher, and R.L. Sass. 1997. Comparison of soil acetate concentrations and methane production, transport, and emission in two rice cultivars. *Global Biogeochem. Cycles* 11:1–14. doi:10.1029/96GB03040
- Smith, P., D. Martino, Z. Cai, D. Gwary, H. Janzen, P. Kumar, B. McCarl, S. Ogle, F. O'Mara, C. Rice, B. Scholes, and O. Sirotenko. 2007. 2007: Agriculture. In: B. Metz, O.R. Davidson, P.R. Bosch, R. Dave, and L.A. Meyer, editors, *Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge Univ. Press, Cambridge, UK, and New York. p. 497–540.
- Suralta, R.R., and A. Yamauchi. 2008. Root growth, aerenchyma development, and oxygen transport in rice genotypes subjected to drought and waterlogging. *Environ. Exp. Bot.* 64:75–82. doi:10.1016/j.envexpbot.2008.01.004
- Tyler, S.C., R.S. Bilek, R.L. Sass, and F.M. Fisher. 1997. Methane oxidation and pathways of production in a Texas paddy field deduced from measurements of flux,  $\delta^{13}\text{C}$ , and  $\delta\text{D}$  of  $\text{CH}_4$ . *Global Biogeochem. Cycles* 11:323–348. doi:10.1029/97GB01624
- van Groenigen, J.W., G.L. Velthof, O. Oenema, K.J. van Groenigen, and C. van Kessel. 2010. Towards an agronomic assessment of  $\text{N}_2\text{O}$  emissions: A case study for arable crops. *Eur. J. Soil Sci.* 61:903–913. doi:10.1111/j.1365-2389.2009.01217.x
- Wang, Z.P., C.W. Lindau, R.D. Delaune, and W.H. Patrick, Jr. 1993. Methane emission and entrapment in flooded rice soils as affected by soil properties. *Biol. Fertil. Soils* 16:163–168. doi:10.1007/BF00361401
- Wang, B., H.U. Neue, and H.P. Samonte. 1997. Effect of cultivar difference ('IR72', 'IR65598', 'Dular') on methane emission. *Agric. Ecosyst. Environ.* 62:31–40. doi:10.1016/S0167-8809(96)01115-2
- Wassmann, R., and M.S. Aulakh. 2000. The role of rice plants in regulating mechanisms of methane emissions. *Biol. Fertil. Soils* 31:20–29. doi:10.1007/s003740050619
- Wassmann, R., M.S. Aulakh, R.S. Lantin, H. Rennenberg, and J.B. Aduna. 2002. Methane emission patterns from rice fields planted to several rice cultivars for nine seasons. *Nutr. Cycling Agroecosyst.* 64:111–124. doi:10.1023/A:1021171303510
- Watanabe, A., M. Kajiura, T. Tashiro, and M. Kimura. 1995. Influence of rice cultivar on methane emission from paddy fields. *Plant Soil* 176:51–56. doi:10.1007/BF00017674
- Xu, H., X. Guangxi, Z. Cai, and H. Tsuruta. 1997. Nitrous oxide emissions from three rice paddy fields in China. *Nutr. Cycling Agroecosyst.* 49:23–28.
- Xu, Y.C., Q.R. Shen, M.L. Li, K. Dittert, and B. Sattelmacher. 2004. Effect of soil water status and mulching on  $\text{N}_2\text{O}$  and  $\text{CH}_4$  emission from lowland rice field in China. *Biol. Fertil. Soils* 39:215–217. doi:10.1007/s00374-003-0692-4
- Yan, X., H. Akiyama, K. Yagi, and H. Akimoto. 2009. Global estimations of the inventory and mitigation potential of methane emissions from rice cultivation conducted using the 2006 Intergovernmental Panel on Climate Change Guidelines. *Global Biogeochem. Cycles* 23:GB2002. doi:10.1029/2008GB003299. doi:10.1029/2008GB003299
- Yan, X., S. Shi, L. Du, and G. Xing. 2000. Pathways of  $\text{N}_2\text{O}$  emission from rice paddy soil. *Soil Biol. Biochem.* 32:437–440. doi:10.1016/S0038-0717(99)00175-3
- Yan, X., K. Yagi, H. Akiyama, and H. Akimoto. 2005. Statistical analysis of the major variables controlling methane emissions from rice fields. *Glob. Change Biol.* 11:1131–1141. doi:10.1111/j.1365-2486.2005.00976.x
- Yu, K.W., Z.P. Wang, and G.X. Chen. 1997. Nitrous oxide and methane transport through rice plants. *Biol. Fertil. Soils* 24:341–343. doi:10.1007/s003740050254
- Zhang, G.B., Y. Ji, J. Ma, G. Liu, H. Xu, and K. Yagi. 2013. Pathway of  $\text{CH}_4$  production, fraction of  $\text{CH}_4$  oxidized, and  $^{13}\text{C}$  isotope fractionation in a straw-incorporated rice field. *Biogeosciences* 10:3375–3389. doi:10.5194/bg-10-3375-2013